arXiv:1209.5976v2 [q-fin.PR] 27 Sep 2012

Quadratic hedging schemes for general GARCH models

Alexandru Badescu[1], Robert J. Elliott[2,3], and Juan-Pablo Ortega[4]

Abstract

In this paper we propose different schemes for option hedging when asset returns are modeled by dynamics from a general class of GARCH models. Since the minimal martingale measure fails to produce a probability measure in this setting, we construct local risk minimization hedging strategies with respect to a risk neutral measure. Local risk minimization is investigated in the context of Gaussian driven models, and two other minimum variance hedges are proposed in order to extend Duan's [31] delta hedge. These two methods are constructed using local risk minimizing hedges for bivariate diffusion limits of GARCH models. Numerical experiments are carried out in order to compare the different hedging schemes; in particular, the sensitivity of the hedging strategies with respect to several pricing measures is tested for a special class of non-Gaussian GARCH models for European style options with different moneyness and maturities.

Keywords: GARCH models, hedging scheme, local risk minimization, conditional Esscher transform, Extended Girsanov Principle, bivariate diffusion limit, minimum variance hedge.

1 Introduction

Many empirical studies have shown strong evidence against some of the underlying assumptions of the Black-Scholes model [12], in particular the constant value that is assumed for the volatility. The incorporation of a stochastic volatility in modelling the dynamics of asset returns plays a major role in explaining some of the stylized properties of financial time series. Consequently, the pricing and hedging of options based on stochastic volatility (SV) models have been frequently studied in the mathematical finance literature in both discrete and continuous time. Discrete-time discussions have been dominated by the Generalized Autoregressive Conditionally Heteroskedastic (GARCH) models introduced by Engle [41] and Bollerslev [13].

The majority of the GARCH option pricing literature relies on the locally risk neutralized valuation relationship (LRNVR) proposed by Duan [31]. This principle is constructed using an equilibrium argument and generalizes the risk-neutral valuation methodology first proposed by Rubinstein [65] and later developed by Brennan [15]. The key assumption behind the construction of the pricing kernel is the conditionally bivariate Gaussian distribution of the

---

[1] Department of Mathematics and Statistics, University of Calgary, Calgary, Canada. `abadescu@math.ucalgary.ca`

[2] Department of Mathematics and Statistics, University of Calgary, Calgary, Canada. `relliott@ucalgary.ca`

[3] School of Mathematical Sciences, University of Adelaide, SA 5005, Australia

[4] Centre National de la Recherche Scientifique, Département de Mathématiques de Besançon, Université de Franche-Comté, UFR des Sciences et Techniques. 16, route de Gray. F-25030 Besançon cedex. France. `Juan-Pablo.Ortega@univ-fcomte.fr`

asset returns and changes in the aggregate consumption. Therefore, option pricing using this approach can only be performed within the restrictive family of GARCH models driven by normal innovations. Various ways have been proposed for pricing European style options under such models. The traditional approach is to evaluate the risk neutral expectations of discounted payoffs by combining standard Monte-Carlo simulations with different variance reduction techniques, (see for example, Duan [31], Duan *et al* [37, 36], Christoffersen and Jacobs [21], Hsieh and Ritchken [53] among many others). Another direction has been proposed by Heston and Nandi [51] who obtain semi closed form solutions for European option prices when assets are modeled by affine Gaussian GARCH models.

Although GARCH models provide a better setting than their homoskedastic alternatives, the Gaussian assumption is too restrictive to allow the capture of the evolution of the asset returns and the anomalies observed in the market option quotes. Therefore, different interrelated extensions have been proposed recently in the literature. They can be classified in two main directions. Firstly, new GARCH option pricing models based on a plethora of skewed and leptokurtic non-Gaussian innovations have been suggested; for example, the General Error Distribution in Duan [34], the Inverse Gaussian in Christoffersen *et al.* [23], the Normal Inverse Gaussian in Stentoft [68], the Generalized Hyperbolic in Chorro *et al.* [20], and Gaussian mixtures in Badescu *et al.* [3] or Stentoft and Rombouts [69], among others. Secondly, semi or non-parametric pricing models have also been proposed; see for example Badescu and Kulperger [3] and Barone-Adesi *et al.* [9]. For further recent advances in the GARCH option pricing literature we refer to the survey paper by Christoffersen *et al.* [26] and references therein.

Since markets described by all these discrete-time models are incomplete, a natural question that needs to be answered is the choice of the stochastic discount factor for pricing purposes. Indeed, considering new risk neutral measures is a necessity since the LRNVR cannot be applied outside the Gaussian framework, even though it should be mentioned that Duan provided a generalization of the LRNVR in Duan [34] using an equilibrium argument and invoking conditional bivariate Gaussian assumptions on a function of the asset returns and aggregate consumptions; Badescu *et al.* [5] have shown that this transformation is closely related to a Wang [70] type transform often used for pricing insurance contracts in the actuarial science literature. This pricing kernel has been empirically tested by several studies that include Christoffersen *et al.* [22] and Stentoft [68].

A good candidate among the class of martingale measures is the conditional Esscher transform that was first proposed in a general GARCH setting by Siu *et al.* [67] and has been further investigated in terms of option prices by other authors, such as Badescu and Kulperger [3], and Christoffersen *et al.* [25]. Gourieroux and Monfort [50] also used this exponentially affine stochastic discount factor for a more general class of discrete-time models. Badescu *et al.* [5] provided conditions under which this change of measure is consistent with pricing kernels constructed on consumption CAPM models. Another option concerning the choice of risk neutral measure was proposed by Elliott and Madan [40] as a discrete-time generalization of the Girsanov theorem; this direction has been empirically investigated in Badescu *et al.* [3]. Finally, GARCH option pricing models based on second order conditional Esscher transforms and variance dependent kernels have been recently proposed by Monfort and Pegoraro [60] and Christoffersen *et al.* [24], respectively. Unlike the previously described martingale measures, these require the use of option data, due to the presence of extra parameters which are not estimated based only on historical return data.

Despite their popularity in terms of option pricing, GARCH models have rarely been used for option hedging purposes. The first such proposal can be found in the paper by Duan [31] which introduced a delta hedge obtained as the partial derivative of the option price with respect to underlying and computes it using Monte Carlo simulation for evaluating the corresponding LRNVR expectation. Other approaches for computing this hedge ratio include the finite difference method proposed by Engle and Rosenberg [42] or the adjusted Black Scholes delta hedging computations based on analytic approximations of GARCH option prices (see Choi [27]). Kallsen and Taqqu [58] proposed a continuous time GARCH process under which the associated markets are complete and also investigated hedging in the Black-Scholes sense. Duan's [31] definition of the delta hedge has been subjected to several criticisms, the most important coming from Garcia and Renault [48] who pointed out that this specification ignores the fact that the option price depends also on the conditional variance, which is itself a function of the asset price. In order to take into account the stochastic volatility environment of GARCH models, Engle and Rosenberg [43] considered the computation of the option's deltas and gammas using the chain rule and studied the modifications in empirical performance when hedging options with changing variance term structure for different volatility specifications.

The techniques briefly described above try to extend the hedging designed for complete markets to a GARCH context, and therefore they are not optimal in the incomplete market setting. Moreover, the use of continuous time hedging ratios for discrete-time models may generate high hedging errors. Thus, various other hedging techniques have been proposed in the financial literature by minimizing different measures for the hedging error, such as quadratic functionals. One of the most influential strategies in this direction has been the local risk minimization scheme introduced by Föllmer, Schweizer, and Sondermann (see [45, 47, 66], and references therein); this approach uses the notion of sequential regressions in order to construct a generalized trading strategy (not necessarily self-financing) that minimizes daily squared hedging errors. Local risk minimization is particularly convenient in the discrete time setup because it is well adapted to recursive algorithmic implementations and can be tuned to minimize hedging errors related to arbitrary prescribed hedging frequencies. The standard approach for constructing local risk minimization hedges is to minimize the conditional expected value of the quadratic cost of hedging under the historical probability measure. This optimization procedure also gives rise to a contingent claim price computed in the risk neutral world with the well-known minimal martingale measure. Although local risk minimization has been widely used for pricing and hedging financial derivatives based on continuous time models, its implementation in discrete-time, and in particular for GARCH models, raises some issues. For example, in Ortega [62] it has been shown that in the GARCH context it is only under very restrictive hypotheses (the model innovations are required to be bounded) that option prices arising from local risk minimization using the physical measure are guaranteed to be arbitrage free. Ortega also shows that in the absence of those hypotheses, the minimal martingale measure, whose existence ensures that the local risk minimizing price is free of arbitrage, is signed. Additionally, a number of situations have been identified (see e.g. Černý and Kallsen [17]) in which the local minimization of squared hedging errors does not guarantee the minimization of the final unconditional expected squared hedging error; this point has motivated recent important work which target the minimization of the global hedging risk (see e.g. Černý and Kallsen [17, 18, 19]). Therefore, in order to address some of these issues, quadratic hedging with respect to a martingale measure has been recently proposed. When the local risk minimization is carried out with respect to a martingale measure, a number of interesting features arise. For example, the option prices that result from the scheme are guaranteed to be arbitrage free and the hedging strategies in this case minimize, not only the local but also the global hedging risk. Additionally, there is a decoupling between the expressions that determine hedges and prices that can be taken advantage of in order to create numerically performing schemes to compute them. Additional advantages of using this approach are discussed in Section 3.1; an intuitive explanation behind risk neutral optimization and references to other research studies which undertake this approach are presented in Cont *et al.* [28]. To our knowledge, the only attempts at exploring local risk minimization in a GARCH setting are due to Ortega [62] and Huang and Guo [55]. The former investigates the use of the local risk minimization with respect to a martingale measure in the context of Gaussian driven GARCH processes and shows how Duan [31] and Heston and Nandi's [51] pricing formulas can be recovered in this fashion, while the latter investigates the risk adjusted expected quadratic cost of hedging minimal hedging strategies in the sense of Elliott and Madan [40]. The empirical findings of these studies show that such techniques provide a viable alternative to the standard hedge ratios used for GARCH models.

In this paper we provide a unified framework for the local risk minimization and the standard continuous-time delta hedges for a general class of GARCH models. Firstly, we introduce local risk minimization with respect to a martingale measure for different hedging frequencies under a GARCH model with an unspecified distribution for the driving noise. We present the main advantages of using such a scheme. Although this construction is valid for any risk neutral measure, we shall focus on the conditional Esscher transform and on the Extended Girsanov Principle for our numerical examples. These two measures are identical and coincide with the one coming from Duan's LRNVR [31] for normally driven GARCH processes, so it is only in the case of non-Gaussian driven models where the effect of the risk neutral measure on the hedging performance will be investigated.

When the time between two consecutive observations approaches zero, we show that both the local risk minimization hedge ratio under a martingale measure and the standard one-step ahead minimum variance GARCH hedge taken with respect to the historical measure can be approximated by a stochastic volatility type delta hedge computed using a chain rule differentiation for the total derivative. More specifically, the resulting delta consists of the sum of the option delta (the first partial derivative of the option price with respect to the spot price) and the option vega (the first partial derivative of the option price with respect to the conditional variance process) scaled by a so-called vega multiplier (VM). The name of this term is borrowed from Engle and Rosenberg [43]. However, this approximation is different from the one proposed by Engle and Rosenberg [43] since all our hedge ratios and correction terms are computed directly based on a GARCH model instead of on an adjusted Black-Scholes price in the sense of Hull and White [56]. Further derivations for the GARCH delta and vega are computed in the case of a Gaussian model with a generalized volatility dynamic, but these quantities require Monte Carlo simulation for their evaluation. For a particular case of a Gaussian asymmetric GARCH model which contains a leverage effect, we derive the corresponding local risk minimization hedge of its continuous time bivariate diffusion limit taken in the sense of Duan [33]. This result may also serve as a further justification of why Duan's [31] hedge is the correct instantaneous delta hedge only when changes in today's return and tomorrow's variance are conditionally uncorrelated. However, for the levered Gaussian GARCH model, the corresponding VM is non-zero. Thus an adjustment to Duan's delta hedge has to be made. In this direction, we propose two alternative hedging schemes: the first one is based on the continuous time limit of the GARCH VM and the second uses the standard chain rule differentiation, so the VM is calculated as the derivative of the variance process with respect to the asset price. Our numerical experiments suggest that the local risk minimization and the delta hedge computed based on the continuous time approximations outperform the other hedging scheme considered for almost all maturities, moneyness and hedging frequencies considered. The performance measure used for our comparison purposes is based on the mean square hedging error (MSHE) constructed from replicating European call options. Next, we compare the hedging performance of the local risk minimization and Duan's delta hedge when the returns are modeled with a Normal Inverse Gaussian GARCH model under the conditional Esscher transform and the Extended Girsanov Principle. Similar numerical results suggest that there are no significant differences between the hedging schemes when these two pricing measures are used. Additionally, our findings suggest that overall, the local risk minimization taken with respect to a martingale measure outperforms Duan's delta hedge for at-the-money and out-of-money contracts and in the cases when the portfolio rebalancing is made less frequently.

The paper is organized as follows. Section 2 characterizes the risk neutral GARCH dynamics for the pricing kernel used. Section 3 presents the local risk minimization and its relationship with the continuous time hedging schemes. In Section 4 we explicitly compute the delta and the vega hedges for GARCH models and show the relationship between local risk minimization under GARCH and its continuous time limit. Numerical experiments that compare the performances of the different hedging schemes are provided. Section 5 studies different implementations of the local risk minimization hedging scheme for models with Normal Inverse Gaussian innovations using the martingale measures which were introduced in Section 2. Section 6 concludes the paper.

2 GARCH(1,1) models and martingale measures

Consider a discrete time financial market with the set of trading dates indexed by $\mathcal{T} = \{t | t = 0, \ldots, T\}$. The market consists of one reference and one risky asset. We denote by P the underlying probability measure and we assume that the dynamics of these assets are modeled by the following bivariate stochastic process $(S^0, S) = \{(S^0_t, S_t)\}_{t\in\mathcal{T}}$. The risk-free asset is a deterministic process and it evolves over time according to the following equation:

$$S^0_t = e^{rt}, \quad S^0_0 = 1. \tag{2.1}$$

Here r represents the constant one-period continuously compounded risk free rate of return. We also denote by $\widetilde{S}_t$ the discounted asset price process from time $\tau = t$ to $\tau = 0$, so $\widetilde{S}_t = e^{-rt}S_t$.

The log-return on the risky asset process, denoted by $y = \{y_t\}_{t\in\mathcal{T}} := \{\ln(S_t/S_{t-1})\}_{t\in\mathcal{T}}$, has a general GARCH(1,1) structure with the P-dynamics given below:

$$\begin{aligned} y_t &= \mu_t + \sigma_t\epsilon_t, \quad \epsilon_t|\mathcal{F}_{t-1} \sim \mathbf{D}(0,1), & (2.2)\\ \sigma_t^2 &= \alpha_0 + \alpha_1\sigma_{t-1}^2\omega(\epsilon_{t-1}) + \beta_1\sigma_{t-1}^2. & (2.3)\end{aligned}$$

The following assumptions and notations complete our description under P:

- The uncertainty modeled through the complete probability space $(\Omega, \mathcal{F}, P)$ has a filtration $\mathcal{F} := \{\mathcal{F}_t\}_{t\in\mathcal{T}}$, where $\mathcal{F}_t$ represents the σ-field of all market information available up to time t.

- $\mathcal{F}_t = \sigma(\epsilon_1, \ldots, \epsilon_t)$ where $\{\epsilon_t\}_{t\in\mathcal{T}}$ is a sequence of random variables such that for each t, ϵ_t is independent and identically distributed conditional on $\mathcal{F}_t$ with a standard distribution $D(0,1)$ under P; the two numbers in the symbol $D(0,1)$ indicate that this distribution has mean 0 and variance 1. We denote by $f^P_{\epsilon_t}(\cdot)$ and $F^P_{\epsilon_t}(\cdot)$ the $\mathcal{F}_{t-1}$-conditional probability density function and cumulative distribution function of ϵ_t under P, respectively; we further assume that its conditional cumulant generating function exists in a neighborhood of zero:
$$\kappa^P_{\epsilon_t}(z) = \ln E^P_{t-1}[e^{z\epsilon_t}] < \infty, \quad z \in (-u, u), \ \ u > 0.$$

- The conditional mean log-return μ_t is $\mathcal{F}_{t-1}$ measurable. Different mean specifications, similar to the ones proposed in Duan [31] or Christoffersen *et al* [25], are discussed in the following sections after parametric assumptions for the driving noise are made.

- The conditional variance equation has a GARCH(1,1) structure with the news impact curve modeled through the function $\omega(\cdot)$. The GARCH parameters are assumed to satisfy the standard second order stationarity conditions.

The following lemma summarizes two results later used in this paper for constructing locally risk minimizing hedging strategies under P. The proof involves basic algebraic manipulations of conditional moments of (2.2) and is therefore omitted.

Lemma 2.1 *The physical mean and variance of changes in discounted asset prices are given by:*

$$\begin{aligned} E^P\left[\widetilde{S}_t - \widetilde{S}_{t-1} \mid \mathcal{F}_{t-1}\right] &= e^{\mu_t - rt} S_{t-1}\left(e^{\kappa^P_{\epsilon_t}(\sigma_t)} - e^{r-\mu_t}\right), & (2.4)\\ \mathrm{Var}^P\left[\widetilde{S}_t - \widetilde{S}_{t-1} \mid \mathcal{F}_{t-1}\right] &= e^{2(\mu_t - rt)} S^2_{t-1}\left[e^{\kappa^P_{\epsilon_t}(2\sigma_t)} - e^{2\kappa^P_{\epsilon_t}(\sigma_t)}\right]. & (2.5)\end{aligned}$$

It is well known that, in the absence of arbitrage, the price of any contingent claim can be expressed as the discounted expected value of its payoff at maturity under a P - equivalent risk neutral probability measure. The discrete-time structure of our model makes the market incomplete and, therefore, there exist an infinite number of such pricing measures. In order to complete the characterization of our GARCH(1,1) pricing model we need to specify the risk neutralized dynamic of the log-returns under such a measure. Thus, we propose below the following general GARCH(1,1)-type structure for the log-return process under a probability martingale measure generically denoted by Q:

$$\begin{aligned} y_t &= r - \kappa^Q_{\epsilon^*_t}(\sigma^*_t) + \sigma^*_t\epsilon^*_t, \quad \epsilon^*_t|\mathcal{F}_{t-1} \sim \mathbf{D}^*(0,1) & (2.6)\\ \sigma^{*2}_t &= \alpha^*_{0t} + \alpha^*_{1t}\sigma^{*2}_{t-1}\omega^*(\epsilon^*_{t-1}) + \beta^*_{1t}\sigma^{*2}_{t-1}. & (2.7)\end{aligned}$$

The above equations require some explanation and further comments as follows:

- All risk neutral quantities are illustrated through a "*" notation, so that σ_t^{*2} represents the risk neutral conditional variance of the log-return process, while $\kappa^Q_{\epsilon^*_t}(\cdot)$ is the $\mathcal{F}_{t-1}$- conditional cumulant generating function of ϵ^*_t under Q.

- The GARCH - type parameters are no longer constants under the new measure. Instead, in order to keep the predictability structure of the conditional variance under Q, we assume that α^*_{0t}, α^*_{1t} and β^*_{1t} are all $\mathcal{F}_{t-1}$ measurable.

- The standard innovation process under the martingale measure is assumed to have a conditional distribution under the martingale measure $\mathbf{D}^*(0,1)$ different from $\mathbf{D}(0,1)$. However, in many practical examples it is often the case that $\mathbf{D}^*$ and $\mathbf{D}$ coincide.

We can now formulate the results from Lemma 2.1 in the risk neutral case. The proof follows immediately by replacing the risk neutral return specifications into Lemma 2.1 .

Lemma 2.2 *The risk neutral mean and variance of changes in discounted asset prices are given by:*

$$\begin{aligned} E^Q\left[\widetilde{S}_t - \widetilde{S}_{t-1} \mid \mathcal{F}_{t-1}\right] &= 0, & (2.8)\\ \mathrm{Var}^Q\left[\widetilde{S}_t - \widetilde{S}_{t-1} \mid \mathcal{F}_{t-1}\right] &= e^{-2r(t-1)} S^2_{t-1}\left[e^{\kappa^Q_{\epsilon^*_t}(2\sigma^*_t) - 2\kappa^Q_{\epsilon^*_t}(\sigma^*_t)} - 1\right]. & (2.9) \end{aligned}$$

Note that equation (2.8) is a direct consequence of the fact that discounted asset prices are martingales under Q. The dynamics from (2.6)-(2.7) provide us with a very general GARCH characterization in the risk neutral world. This structure nests some of the most popular martingale measures used in the GARCH option pricing literature. In the remainder of this section we show that both the conditional Esscher transform and the Extended Girsanov Principle give rise to risk neutral dynamics which are special cases of our proposed model.

2.1 Extended Girsanov Principle

An extended Girsanov principle has been proposed by Elliott and Madan [40] as an alternative pricing measure for discrete time models. As its name suggests, the construction of this measure is based on an extension of the Girsanov transformation to non-Gaussian settings according to which asset prices will have a different mean return and an unchanged volatility after the change of measure. This measure has been used for option pricing by Badescu and Kulperger [3] in a semiparametric GARCH setting and Badescu *et al* [4] for Gaussian mixture GARCH models.

Although the Radon-Nikodym derivative is defined in terms of the martingale component process from the multiplicative Doob decomposition of the discounted asset price, we follow Badescu *et al* [6] and characterize the risk neutralized dynamic for our proposed generalized GARCH setup. The resulting equations can be viewed as a special case of (2.6)-(2.7).

Proposition 2.3 *Define the stochastic process* $N := \{N_t\}_{t\in\mathcal{T}}$*:*

$$N_t = \frac{f^P_t(\epsilon_t + \varrho_t)}{f^P_t(\epsilon_t)},$$

Here $\{\varrho_t\}_{t\in\mathcal{T}}$ is an $\mathcal{F}_t$-predictable process representing the market price of risk given by:

$$\varrho_t = \frac{\mu_t + \kappa^P_{\epsilon_t}(\sigma_t) - r}{\sigma_t}. \tag{2.10}$$

Then, the following statements hold:

(i) *The process $Z := \{Z_t\}_{t\in\mathcal{T}}$ defined by $Z_t := \prod_{k=1}^{t} N_k$ with $Z_0 = 1$ is a P-martingale and Z_T defines an equivalent measure Q by $Z_T = \frac{dQ}{dP}$.*

(ii) *The innovation process $\epsilon^* := \{\epsilon^*_t\}_{t\in\mathcal{T}}$ with $\epsilon^*_t = \epsilon_t + \varrho_t$ is a sequence of $\mathcal{F}_{t-1}$-conditionally uncorrelated zero mean and unit variance random variables under Q.*

(iii) *The risk neutralized asset return dynamic is given by:*

$$\begin{aligned} y_t &= r - \kappa^Q_{\epsilon^*_t}(\sigma_t) + \sigma_t\epsilon^*_t, \quad \epsilon^*_t \sim \mathbf{D}(0,1), \\ \sigma^2_t &= \alpha_0 + \alpha_1\sigma^2_{t-1}\omega(\epsilon^*_{t-1} - \varrho_{t-1}) + \beta_1\sigma^2_{t-1}. \end{aligned}$$

In order to differentiate among the risk neutral measures used throughout this paper, we further denote the measure associated to the Extended Girsanov transform by Q_{egp}. For completeness, the proof of this proposition is provided in the Appendix.

It is not difficult to see that the above dynamic is a particular case of (2.6)-(2.7) when the risk neutral innovation is shifted by the risk premium process and the conditional variance under P remains unchanged after the measure change (i.e. $\sigma_t = \sigma^*_t$). Although in our numerical simulations we use special cases of this result, this risk neutralized dynamic can be applied for pricing and hedging under GARCH models for any distribution which satisfy the constraints imposed of the beginning of the section.

Remark 2.4 An important characteristic of the Extended Girsanov Principle is its relationship to the notion of locally risk minimizing hedging. Elliott and Madan [40] showed that this new martingale measure is the unique probability measure which is consistent with hedging strategies that minimize the one-step ahead quadratic risk adjusted cost of hedging. Further comments on such strategies are made at the end of Section 3.1 after we introduce the notion of locally risk minimizing strategies.

2.2 Conditional Esscher transform

The conditional Esscher transform has been first introduced into a GARCH setting by Siu *et al.* [67]. Since then, this transformation has been used in a variety of empirical studies which dealt with the pricing of financial derivatives under different assumptions regarding the innovation process. For example, Badescu *et al.* [4] and Stentoft and Rombouts [69] used it for pricing both European and American style options under Gaussian mixture innovations. This method has been also investigated and further generalized by Christoffersen *et al.* [25]. The change of measure is based on an exponential affine stochastic discount factor and it has become a very popular tool for many of the GARCH option pricing models due to its Radon-Nikodym derivative tractability. Another appealing feature of this change of measure is the fact that it preserves the parametric driving noise distribution family after the change of measure. For example, Siu *et al* [67] and Christoffersen *et al* [25] derived such a result for infinite divisible distributions.

In the following proposition we characterize the risk neutralized dynamic of the asset return process given in (2.2)-(2.3) under the conditional Esscher transform and we show this is a special case of the general family of GARCH pricing models from (2.6)-(2.7).

Proposition 2.5 *Define the stochastic process* $N := \{N_t\}_{t\in\mathcal{T}}$*:*

$$N_t = e^{\theta_t\sigma_t\epsilon_t - \kappa^P_{\epsilon_t}(\theta_t\sigma_t)},$$

Here $\{\theta_t\}_{t\in\mathcal{T}}$ *is a* $\mathcal{F}_t$*-predictable process which satisfies the following martingale condition:*

$$\mu_t + \kappa^P_{\epsilon_t}\big((1+\theta_t)\sigma_t\big) - \kappa^P_{\epsilon_t}\big(\theta_t\sigma_t\big) = r. \tag{2.11}$$

Then, the following statements hold:

(i) *The process* $Z := \{Z_t\}_{t\in\mathcal{T}}$ *defined by* $Z_t := \prod_{k=1}^{t} N_k$ *with* $Z_0 = 1$ *is a* P*-martingale and* Z_T *defines an equivalent measure* Q *by* $Z_T = \frac{dQ}{dP}$*.*

(ii) *The innovation process* $\epsilon^* := \{\epsilon^*_t\}_{t\in\mathcal{T}}$ *defined by:*

$$\epsilon^*_t = \frac{\epsilon_t - \kappa^{P\prime}_{\epsilon_t}(\theta_t\sigma_t)}{\sqrt{\kappa^{P\prime\prime}_{\epsilon_t}(\theta_t\sigma_t)}},$$

is a sequence of $\mathcal{F}_{t-1}$*-conditionally uncorrelated zero mean and unit variance random variables under* Q*. Here* $\kappa^{P\prime}$ *and* $\kappa^{P\prime\prime}$ *represent the first and second order derivatives of the corresponding cumulant generating functions.*

(iii) *The risk neutralized asset return dynamic is given by:*

$$\begin{aligned} y_t &= r - \kappa^Q_{\epsilon^*_t}(\sigma^*_t) + \sigma^*_t\epsilon^*_t, \quad \epsilon^*_t \sim \mathbf{D}^*(0,1), \\ \sigma^{*2}_t &= \alpha^*_{0t} + \alpha^*_{1t}\sigma^{*2}_{t-1}\omega(\kappa^{P\prime}_{\epsilon_t}(\theta_t\sigma_t) + \sqrt{\kappa^{P\prime\prime}_{\epsilon_t}(\theta_t\sigma_t)}\epsilon^*_{t-1}) + \beta^*_{1t}\sigma^{*2}_{t-1}. \end{aligned}$$

Here the conditional variance dynamic coefficients are given by:

$$\alpha^*_{0t} = \alpha_0\kappa^{P\prime\prime}_{\epsilon_t}(\theta_t\sigma_t), \quad \alpha^*_{1t} = \alpha_1\frac{\kappa^{P\prime\prime}_{\epsilon_t}(\theta_t\sigma_t)}{\kappa^{P\prime\prime}_{\epsilon_{t-1}}(\theta_{t-1}\sigma_{t-1})}, \quad \beta^*_{1t} = \beta_1\frac{\kappa^{P\prime\prime}_{\epsilon_t}(\theta_t\sigma_t)}{\kappa^{P\prime\prime}_{\epsilon_{t-1}}(\theta_{t-1}\sigma_{t-1})}.$$

Although some proofs of parts of the above theorems are already given in a less general context in some other papers, for completeness we provide a complete proof of the theorem in the Appendix. We denote by Q_{ess} the change of measure induced by the conditional Esscher transform. It is easy to note that the above risk neutral specification is again a special case of the general risk-neutralized GARCH model introduced at the beginning of the section. Unlike the Extended Girsanov Principle, the conditional Esscher transform gives rise to a different risk neutral conditional variance. For example, using a second order Taylor expansion for the second order derivative of the cumulant generating function $\kappa^P_{\epsilon_t}(\theta_t \sigma_t)$ around zero, Christoffersen *et al* [25] show that the risk neutral conditional variance will be always larger than its physical counterpart provided that the conditional skewness and excess kurtosis are negative and positive, respectively, which is a common stylized fact exhibited by most financial time series. A higher implied volatility, computed from option prices, than the physical volatility is a well-known empirical finding in the option pricing literature and gives rise to the necessity of incorporating a volatility risk premium. Note that the risk neutral dynamic no longer has a standard GARCH structure since the parameters α^*_{0t}, α^*_{1t} and β^*_{1t} are $\mathcal{F}_t$ predictable processes, which makes it a special case of the proposed GARCH risk neutral equations (2.6)-(2.7).

A second implication of the above result is that Esscher transforms does not preserve the innovation distribution under the change of measure (i.e. $\mathbf{D}^* \neq \mathbf{D}$). However, this is not be the case for our numerical simulations which are based on Gaussian and Normal Inverse Gaussian (NIG) GARCH models.

Finally, the process θ_t is a unique solution of the martingale condition in **(ii)**, which uniquely characterizes the Esscher transform. An explicit solution of (2.11) is rarely available and solving numerically for θ_t can be computationally demanding for an option pricing exercise, thus making the implementation of the local risk minimization hedging scheme not feasible. An alternative solution is to use a Taylor expansion around the origin for both cumulant generating functions in (2.11) in order to obtain an approximation. Such a strategy leads to the following approximate solution:

$$\hat{\theta}_t \approx \frac{r - \mu_t}{\sigma_t^2} - \frac{1}{2}. \tag{2.12}$$

The above approximation coincides with the exact solution when innovations are Gaussian and in this case the risk neutral variance equals the physical one. The advantage of using NIG innovations is twofold: firstly, we specify a skewed and leptokurtic density which fits the returns better than the Gaussian one and secondly, the conditional risk neutral variance under the Esscher transform is different from that given by the Extended Girsanov Principle. Moreover, in this setup we have an explicit solution for θ_t which we provide in Section 5.

3 Risk minimization for GARCH options

The market associated with GARCH modeled assets is intrinsically incomplete, that is, there exist contingent products H that cannot be fully replicated using a self-financing portfolio made out of its underlying and a bond. A number of techniques have been developed over the years that tackle simultaneously the pricing and hedging problems by replacing the concept of replication with that of hedging efficiency. As we shall see, these techniques provide simultaneous expressions for prices and hedging ratios which, even though in most cases they require Monte Carlo computations, admit convenient interpretations based on the notion of hedging error minimization that can be adapted to various models for the underlying asset.

In order to provide a brief account of the hedging techniques that we shall be using we start by introducing several definitions.

As we have done so far, we will denote by S_t the price of the underlying asset at time t. The symbol r denotes the constant and continuously composed risk-free interest rate paid on the currency of the underlying in the period that goes from an arbitrary time step $t-1$ to the next t. The price at time t of the riskless asset S^0 such that $S^0_0 = 1$, is hence given by $S^0_t = \mathrm{e}^{rt}$.

Let now $H(S_T)$ be a European contingent claim that depends on the terminal value of the risky asset S_t. As we already pointed out, in the context of an incomplete market, it will be in general impossible to replicate the payoff H by using a self-financing portfolio. Therefore, we introduce the notion of generalized trading strategy, in which the possibility of additional investment in the riskless asset throughout the trading periods up to expiry time T is allowed. All the following statements are made with respect to a fixed filtered probability space $(\Omega, P, \mathcal{F}, \{\mathcal{F}_t\}_{t\in\{0,\ldots,T\}})$, with P being this time around not the physical probability but an arbitrary probability measure.

- A generalized trading strategy is a pair of stochastic processes (ξ^0, ξ) such that $\{\xi^0_t\}_{t\in\{0,\ldots,T\}}$ is adapted and $\{\xi_t\}_{t\in\{1,\ldots,T\}}$ is predictable. The associated value process V of (ξ^0, ξ) is defined as

$$V_0 := \xi^0_0, \quad \text{and} \quad V_t := \xi^0_t \cdot S^0_t + \xi_t \cdot S_t, \quad t \geq 1.$$

- The gains process G of the generalized trading strategy (ξ^0, ξ) is given by

$$G_0 := 0 \quad \text{and} \quad G_t := \sum_{k=1}^{t} \xi_k \cdot (S_k - S_{k-1}), \quad t = 0, \ldots, T.$$

- The cost process C is defined by the difference

$$C_t := V_t - G_t, \quad t = 0, \ldots, T.$$

These processes have discounted versions $\widetilde{V}_t$, $\widetilde{G}_t$, and $\widetilde{C}_t$ defined as:

$$\widetilde{S}_t := S_t \mathrm{e}^{-rt}, \qquad \widetilde{V}_t := V_t \mathrm{e}^{-rt}, \qquad \widetilde{G}_t := \sum_{k=1}^{t} \xi_k \cdot (\widetilde{S}_k - \widetilde{S}_{k-1}), \quad \text{and} \quad \widetilde{C}_t := \widetilde{V}_t - \widetilde{G}_t.$$

Assume now that both H and the $\{S_t\}_{t\in\{0,\ldots,T\}}$ are in $L^2(\Omega, P)$. A generalized trading strategy is called admissible for H whenever it is in $L^2(\Omega, P)$ and its associated value process is such that

$$V_T = H, \quad \text{P } a.s., \qquad V_t, G_t \in L^2(\Omega, P), \quad \text{for each } t.$$

In the context of quadratic criteria, hedging efficiency has been mainly handled in the literature by finding strategies $(\widehat{\xi}^0, \widehat{\xi})$ that minimize one of the three following risk processes:

(i) Local risk process: $LR_t(\xi^0, \xi) := E^P\left[(\widetilde{C}_{t+1} - \widetilde{C}_t)^2 \mid \mathcal{F}_t\right]$, $t = 0, \ldots, T-1$.

(ii) Remaining conditional risk: $RR_t(\xi^0, \xi) := E^P[(\widetilde{C}_T - \widetilde{C}_t)^2 \mid \mathcal{F}_t] = E^P[(H - \sum_{k=t+1}^{T} \xi_k \cdot (\widetilde{S}_k - \widetilde{S}_{k-1}) - V_t)^2 \mid \mathcal{F}_t]$, $t = 0, \ldots, T-1$.

(iii) Global risk: $GR(\xi^0, \xi) := E^P[(\widetilde{C}_T - \widetilde{C}_0)^2] = E^P[(H - \sum_{k=1}^{T} \xi_k \cdot (\widetilde{S}_k - \widetilde{S}_{k-1}) - V_0)^2]$.

We summarize below several reasons why the local risk minimization has been probably the most influential scheme in the literature so far:

- Availability of a convenient characterization of local risk-minimizing strategies and of an easy to implement recursive algorithm to compute them: indeed, it can be shown that an admissible strategy is locally risk-minimizing if and only if the discounted cost process is a P-martingale and it is strongly orthogonal to $\widetilde{S}$, in the sense that $\mathrm{Cov}^P(\widetilde{S}_{t+1} - \widetilde{S}_t, \widetilde{C}_{t+1} - \widetilde{C}_t \mid \mathcal{F}_t) = 0$, P-a.s., for any $t = 0, \ldots, T-1$. Moreover, whenever a local risk-minimizing technique exists, it is fully determined by the backwards recursions:

  $$\begin{aligned} V_T &= H, && (3.1)\\ \xi_{t+1} &= \frac{\mathrm{Cov}^P(\widetilde{V}_{t+1}, \widetilde{S}_{t+1} - \widetilde{S}_t \mid \mathcal{F}_t)}{\mathrm{Var}^P[\widetilde{S}_{t+1} - \widetilde{S}_t \mid \mathcal{F}_t]} = \frac{E^P\left[(\widetilde{V}_{t+1} - \widetilde{V}_t)(\widetilde{S}_{t+1} - \widetilde{S}_t) \mid \mathcal{F}_t\right]}{E^P\left[\widetilde{S}_{t+1} - \widetilde{S}_t \mid \mathcal{F}_t\right]}, && (3.2)\\ \widetilde{V}_t &= E^P\left[\widetilde{V}_{t+1} \mid \mathcal{F}_t\right] - \xi_{t+1} E^P\left[\widetilde{S}_{t+1} - \widetilde{S}_t \mid \mathcal{F}_t\right]. && (3.3) \end{aligned}$$

  The initial investment V_0 determined by these relations is referred to as the local risk minimization option price associated to the measure P.

- The minimization of the local risk implies in many situations the minimization of the global and the remaining risks: for example when P is a martingale measure, local risk minimizing strategies are variance optimal (that is, they minimize the global risk; see [44, Proposition 10.37])) and minimize also the remaining conditional risk (see [44, Proposition 10.34]); ; this is in general not true outside the martingale framework (see [66, Proposition 3.1] for a counterexample). Outside the martingale context things are more complicated and there are in general only sufficient conditions that allow the construction of global risk minimizing strategies out of local ones; these hypothesis are generically not satisfied in the GARCH context; see for example [44, Theorem 10.40].

- Local risk minimization can be adapted to prescribed changes in the hedging frequency: suppose that the life of the option H with maturity in T time steps is partitioned into identical time intervals of duration j; this assumption implies the existence of an integer k such that $kj = T$. We now wish to set up a local risk minimizing replication strategy for H in which hedging is carried out once every j time steps. We shall denote by ξ_{t+j} the hedging ratio at time t that presupposes that the next hedging will take place at time $t+j$. The value of such ratios will be obtained by minimizing the j-spaced local risk process:

  $$LR_t^{(j)}(\xi^0, \xi) := E^P\left[(\widetilde{C}_{t+j}^{(j)} - \widetilde{C}_t^{(j)})^2 \mid \mathcal{F}_t\right], \quad t = 0, j, 2j, \ldots, (k-1)j = T - j. \qquad (3.4)$$

  Here $\{\widetilde{C}_t^{(j)}\}$ is a cost process constructed out of value and gains processes, $\{V_t^{(j)}\}$ and $\left\{G_t^{(j)}\right\}$ that only take into account the prices of the underlying assets at time steps $t = 0, j, 2j, \ldots, kj = T$, in particular, given an integer l such that $t = lj$

  $$\widetilde{G}_t^{(j)} := \sum_{r=1}^{l} \xi_{rj} \cdot (\widetilde{S}_{rj} - \widetilde{S}_{(r-1)j}).$$

  The solution of the minimization problem for (3.4) is the same as (3.1)–(3.3) with the symbols $t+1$ replaced by $t+j$ and the variable t taking the values $t = 0, j, 2j, \ldots, (k-1)j = T-j$. A convoluted but straightforward induction argument shows that the solution of these recursions is:

$$\begin{aligned} V_T^{(j)} &= H, && (3.5)\\ \xi_{t+j} &= \frac{\mathrm{Cov}^P(e^{-rT}H\cdot N_T\cdot N_{T-j}\cdots N_{t+2j},\widetilde{S}_{t+j}-\widetilde{S}_t\mid\mathcal{F}_t)}{\mathrm{Var}^P[\widetilde{S}_{t+j}-\widetilde{S}_t\mid\mathcal{F}_t]}, && (3.6)\\ \widetilde{V}_t^{(j)} &= E_t^P\left[e^{-rT}H\cdot N_T\cdot N_{T-j}\cdots N_{t+j}\right],\quad t=0,j,2j,\ldots,T-j, && (3.7)\end{aligned}$$

Here, for any $t=j,2j,\ldots,T$, the process N_t is given by:

$$N_t=\frac{1-\gamma_t(\widetilde{S}_t-\widetilde{S}_{t-j})}{E^P\left[1-\gamma_t(\widetilde{S}_t-\widetilde{S}_{t-j})\mid\mathcal{F}_{t-j}\right]}=1-\frac{E^P\left[\widetilde{S}_t-\widetilde{S}_{t-j}\mid\mathcal{F}_{t-j}\right]\left(\widetilde{S}_t-E^P\left[\widetilde{S}_t\mid\mathcal{F}_{t-j}\right]\right)}{\mathrm{Var}^P\left[\widetilde{S}_t-\widetilde{S}_{t-j}\mid\mathcal{F}_{t-j}\right]}, \tag{3.8}$$

where,

$$\gamma_t=\frac{E^P\left[\widetilde{S}_t-\widetilde{S}_{t-j}\mid\mathcal{F}_{t-j}\right]}{E^P\left[(\widetilde{S}_t-\widetilde{S}_{t-j})^2\mid\mathcal{F}_{t-j}\right]}.$$

When the product $Z_T:=\prod_{t\in\{j,2j,\ldots,T\}}N_t$ is a positive random variable, the relation $Z_T:=dP/dQ_{\min}$ defines a measure equivalent to P, called the minimal martingale measure, for which the discounted partial process $\{\widetilde{S}_j,\widetilde{S}_{2j},\ldots,\widetilde{S}_T\}$ is a martingale. The interest of this measure is that, when it exists and in the case $j=1$, the value process of the local risk-minimizing strategy *with respect to the physical measure* coincides with arbitrage free price for H obtained by using $Q_{\min}$ as a pricing kernel. Indeed, (3.6) can be rewritten as

$$V_t^{(j)}=E^P\left[e^{-r(T-t)}H(S_T)\cdot N_T\cdot N_{T-j}\cdots N_{t+j}\mid\mathcal{F}_t\right]=E^{Q_{\min}}\left[e^{-r(T-t)}H(S_T)\mid\mathcal{F}_t\right].$$

In a recent paper, Basak and Chabakauri [10] derive the discrete time mean correcting martingale measure as their hedging neutral measure by solving a standard minimum variance problem with a budget constraint. The optimal ratio from (3.6) corresponds to the optimal hedge per unit asset ratio from Proposition 3 in Basak and Chabakauri [10]. However, their hedging formula is slightly different since it is derived based on a budget constraint, which does not hold in the local risk minimization setting.

Although pricing and hedging using the recursion formulas (3.5)–(3.7) might look straightforward, there are various implementation issues that need to be taken care of when asset returns are modeled by GARCH processes; we analyze in detail some of them in the next section and we propose an alternative hedging scheme.

3.1 Local risk minimization with respect to a martingale measure

The minimal martingale measure is usually a signed measure in discrete time settings, so derivative valuation based on this approach might not be suitable. For example, the Radon-Nikodym process defined in (3.8) can take negative values and this may lead to negative prices when computing the expectations from (3.7) using Monte-Carlo simulation. For asset returns governed by GARCH models, one can impose conditions on the parameter set or the innovations which can be too restrictive and make the estimation more challenging. A sufficient condition in the GARCH context requires the driving noise to be bounded (see for example [62, Proposition 2.6]). However, an unreported numerical simulation study indicates that negative values for N_t happen very rarely for GARCH processes driven by Gaussian innovations. This will no longer be the case if a variety of other heavier tailed densities are used, and one of the objective of the current paper is to explore hedging strategies based on such models.

The recursive derivation of a derivative price according to (3.5) - (3.7) is not an easy task even in settings when the minimal martingale measure is a probability measure. Both the option prices and hedging ratios computations require the use of Monte Carlo simulations under the physical measure P and the standard filtration $\{\mathcal{F}_t\}$ generated by the price process, which in the GARCH setup coincides with the one generated by the driving noise, that is $\mathcal{F}_t = \sigma(\epsilon_1, \ldots, \epsilon_t)$. An inspection of the expressions (3.5)–(3.7) shows that in order to compute them we need first an estimation of the stochastic discount factors $\{N_j, \ldots, N_T\}$. When $j = 1$, these factors can be written down in terms of the conditional cumulant generating function; indeed, using (2.4) and (2.5) in (3.8), we obtain:

$$N_t = 1 - \frac{\left(e^{\kappa^P_{\epsilon_t}(\sigma_t)} - e^{r-\mu_t}\right)\left(e^{\sigma_t \epsilon_t} - e^{\kappa^P_{\epsilon_t}(\sigma_t)}\right)}{\left(e^{\kappa^P_{\epsilon_t}(2\sigma_t)} - e^{2\kappa^P_{\epsilon_t}(\sigma_t)}\right)}.$$

However, when $j > 1$ this estimation is necessarily conducted using Monte Carlo simulations whose results are in turn substituted in the expressions (3.5)–(3.7) that are also evaluated using Monte Carlo. The variance of the resulting estimator is too high and makes the results unacceptable. Moreover, the option prices that result from this technique cannot be interpreted as arbitrage free prices if the hedging frequency is lower than periods at which asset prices are observed. Indeed, in the case $j > 1$, the resulting measure is a martingale measure for the discounted partial process $\{\widetilde{S}_j, \widetilde{S}_{2j}, \ldots, \widetilde{S}_T\}$ but not for the entire discounted price process $\{\widetilde{S}_1, \widetilde{S}_2, \ldots, \widetilde{S}_T\}$ and therefore there is a leeway for arbitrage.

As described in Basak and Chabakauri [10], another potential problem with variance minimizing hedging strategies is their time-inconsistency. These authors argue that such inconsistencies can be avoided by either considering that the market is complete or by assuming a zero risk premium for the asset return. The former does clearly not fit our GARCH setting while the latter is equivalent to the discounted asset price being a martingale under the physical measure; hence, one way to overcome the time-inconsistency problem is to consider hedging directly under a risk neutral measure.

The reasons that we mentioned above lead us to explore the local risk minimization strategy for martingale measures, despite the fact that the historical probability associated with the price process represents a more natural approach from a risk management perspective. More specifically, we shall focus on the martingale measures introduced in Section 2. Additionally, given that the trend terms that separate the physical measure from being a martingale measure are usually very small when dealing with daily or weekly financial returns, it is expected that the inaccuracy committed by using a conveniently chosen martingale measure will be smaller than the numerical error that we would face in the Monte Carlo evaluation of the expressions corresponding to the physical measure; see [62, Proposition 3.2] for an argument in this direction. Thus, following Föllmer and Sondermann [45], we derive recursive pricing and hedging relations using risk minimization under martingale measures. These are obtained in a straightforward manner using expressions (3.5)–(3.7) combined with the martingale hypothesis in its statement that makes the stochastic discount factors (3.8) $\{N_t\}_{t\in\{j,2j,\ldots,T\}}$ automatically all equal to 1.

Proposition 3.1 *Let Q be an equivalent martingale measure for the price process $\{S_t\}$ and let $H(S_T)$ be a European contingent claim that depends on the terminal value of the risky asset S_t. The locally risk minimizing strategy every j time steps with respect to the measure Q is determined by the recursions:*

$$V_T^{(j)} = H, \tag{3.9}$$

$$\xi_{t+j} = \mathrm{e}^{-r(T-t)} \frac{E^Q\left[H(S_T)\left(S_{t+j}\mathrm{e}^{-rj} - S_t\right) \mid \mathcal{F}_t\right]}{\mathrm{Var}^Q\left[S_{t+j}\mathrm{e}^{-rj} - S_t \mid \mathcal{F}_t\right]}, \tag{3.10}$$

$$V_t^{(j)} = E^Q\left[e^{-r(T-t)}H(S_T) \mid \mathcal{F}_t\right], \tag{3.11}$$

for all $t = 0, j, 2j, \ldots, T-j$.

For daily hedging ($j = 1$), the denominator of (3.10) can be explicitly written down using Lemma 2.2, in terms of the $\mathcal{F}_t$-conditional cumulant generating function $\kappa^Q_{\epsilon^*_t}$ of ϵ^*_tunder Q and of the risk neutral conditional variance σ^*_t as

$$\mathrm{Var}^Q\left[e^{-r}S_{t+1} - S_t \mid \mathcal{F}_t\right] = S_t^2\left[e^{\kappa^Q_{\epsilon^*_t}(2\sigma^*_{t+1}) - 2\kappa^Q_{\epsilon^*_t}(\sigma^*_{t+1})} - 1\right].$$

For lower hedging frequencies ($j > 1$) this denominator needs to be evaluated by using the Monte Carlo estimator for the variance, that is,

$$\mathrm{Var}^Q\left[e^{-r}S_{t+1} - S_t \mid \mathcal{F}_t\right] \simeq \frac{1}{N}\sum_{i=1}^{N}\left(S^i_{t+j}\mathrm{e}^{-jr} - S_t\right)^2,$$

where $S^1_{t+j}, \ldots, S^N_{t+j}$ are N realizations of the price process under Q at time $t+j$, using paths that have all the same origin S_t at time t.

The recursion formulas require some explanation. First, local risk minimization under a martingale measure does not directly determine the arbitrage free price as in the historical probability optimization case. The option price is, a priori, established by assuming that the discounted value process is a martingale under Q; (3.11) serves as a verification for this fact. The value process $V_t^{(j)}$ is the same for any j, so it does not depend on the hedging frequency. V_0 represents the necessary initial investment to setup the local minimizing strategy to replicate H. This coincides with the arbitrage free pricing for the payoff H and it strongly depends on the choice of the pricing kernel Q.

Secondly, the hedging ratios in (3.10) can be computed using standard Monte Carlo techniques by simulating asset price paths under the risk neutral measure considered. In cases where simulation under Q is not very convenient, one can compute both the numerator and the denominator of (3.10) by simulating historical paths and make use of the corresponding Radon-Nikodym derivative. More precisely, if the change of measure is defined by, $dQ/dP = \prod_{k=1}^{T} N_k$, then for any $\mathcal{F}_T$ measurable function f and any $t \in \{0, \ldots, T\}$,

$$E^Q\left[f \mid \mathcal{F}_t\right] = E^P\left[f\prod_{k=t+1}^{T} N_k \mid \mathcal{F}_t\right].$$

Note that if Q is chosen to be the minimal martingale measure, the discounted value processes from (3.7) and (3.11) coincides, while their corresponding hedging ratios are different. When Q is the pricing measure given by the Extended Girsanov Principle, the hedging scheme from (3.1) is the same as the one obtained in Elliott and Madan [40] by minimizing the expected discounted risk adjusted squared costs of hedging under the historical measure. A detailed discussion about this is presented in Huang and Guo [54, 55] who constructed numerically efficient algorithms that implement this approach to pricing and hedging. Their method, known as LESRAM (Local Expected Squared Risk Minimizing Hedging), takes advantage of the decoupling between the computation of option values and hedges that one obtains in (3.9)–(3.11) by using a martingale measure, in order to interpret the hedges as the coefficients of a regression of the discounted option prices on the discounted prices of the underlying and the bond. Additionally, they introduced a semiparametric valuation method (DSA, Dynamic Semiparametric Approach) in order to evaluate the explained variable of this regression as accurately as possible.

Finally, the optimal hedge ratios from (3.10) minimize the conditional remaining risk and the global risk, since these criteria are equivalent under a martingale measure. Thus, these strategies are also variance optimal.

The empirical studies conducted in the next sections provide a comparison between the hedging strategy performance of the two risk neutral measures described in Section 2 for GARCH models driven by Gaussian and non-Gaussian innovations. The remainder of this section provides of brief discussion between the relationship between local risk minimization and standard delta neutral hedging ratios from continuous time models.

3.2 Local risk minimization and delta hedging

A particularly attractive feature of risk minimization schemes is that the choice of a specific pricing kernel determines simultaneously, and in a consistent way, prices and associated hedging strategies. In this section we focus on continuous time hedging schemes and illustrate their relationship to the discrete time hedges introduced before. A standard approach is to consider a delta neutral portfolio made out of the underlying and the contingent product itself and identify the quantity which should be invested into the asset price such that the instantaneous variance of the portfolio is minimized. These strategies are called minimum variance hedge ratios and have been investigated in both complete and incomplete settings (see for example Boyle and Emanuel [14], Bakshi *at al.* [7] and Alexander and Nogueira [1] among others).

Weak convergence of hedging strategies has been another topic of interest in the mathematical finance literature. In the context of quadratic hedging, Prigent and Scaillet [64] investigate the weak convergence of local risk minimization strategies to the standard delta hedges given by the partial derivatives of the option price with respect to the underlying. In particular, they showed such a convergence result when the asset price is modeled by a stochastic volatility model driven by uncorrelated Brownian motion. A key ingredient in their derivation is provided by the Markovian structure of the model, which is only the case for a GARCH(1,1) setup. However, such a result is beyond the aim of our paper. Instead, in the following proposition, we characterize the instantaneous minimum variance delta hedge in a GARCH environment, and we show its relationship with the local risk minimization hedge.

Proposition 3.2 *Let H be a European contingent product on the underlying price process $\{S_t\}_{t\in\mathcal{T}}$ that has a GARCH model as generating process. Let Q be the risk neutral measure and $\Pi_t = \Pi(S_t, \sigma^2_{t+1}(S_t)) = E^Q\left[e^{r(T-t)}H \mid \mathcal{F}_t\right]$ be its associated price. Construct a portfolio consisting of a long position on a unit of the derivative product and* *a short one on Δ_t units of the underlying. If the time between two consecutive observations approaches zero, the following relationships hold:*

(i) *The minimum variance delta hedge is approximately given by:*

$$\Delta_t \approx \Delta_t^S + \frac{\mathrm{Cov}^P(S_{t+1} - S_t, \sigma_{t+2}^2 - \sigma_{t+1}^2 \mid \mathcal{F}_t)}{\mathrm{Var}^P[S_{t+1} - S_t \mid \mathcal{F}_t]} \Delta_t^\sigma \approx \Delta_t^S + \frac{d\sigma_{t+1}^2}{dS_t} \Delta_t^\sigma. \tag{3.12}$$

Here $\Delta_t^S = \frac{\partial \Pi_t}{\partial S_t}$ is the standard option delta and $\Delta_t^\sigma = \frac{\partial \Pi_t}{\partial \sigma_{t+1}^2}$ is the vega of the option and the coefficients to Δ_t^σ in both approximations are called the vega multipliers (VM).

(ii) *The approximation (3.12) also holds for the Q local risk minimization hedge ξ_t:*

$$\xi_{t+1} \approx \Delta_t^S + \frac{\mathrm{Cov}^Q(S_{t+1} - S_t, \sigma_{t+2}^2 - \sigma_{t+1}^2 \mid \mathcal{F}_t)}{\mathrm{Var}^Q[S_{t+1} - S_t \mid \mathcal{F}_t]} \Delta_t^\sigma \approx \Delta_t^S + \frac{d\sigma_{t+1}^2}{dS_t} \Delta_t^\sigma. \tag{3.13}$$

Hedging under GARCH models is typically done using the standard definition proposed by Duan [31], according to which the hedge ratio is the partial derivative of the option price with respect to the underlying denoted by Δ_t^S. Garcia and Renault [48] point out that this specification ignores the fact that the option price depends also on the conditional variance which is itself a function of the asset price. The second approximation in (3.12) is a direct consequence of this since it resembles the total derivative of the option price with respect to the asset price. Note that this formula resembles the chain rule formulation used in Engle and Rosenberg [43], but there are two major differences. First, our option delta and vega are computed directly based on the GARCH option price, while Engle and Rosenberg [43] used a Hull and White [56] Black-Scholes plug-in type of formula for the option price. Secondly, their option price is based on the average expected volatility, while ours uses the current value of the volatility as an argument, and thus gives rise to a different specifications of the VM. In the case in which the volatility does not depend on the underlying, the formula reduces to the standard delta hedge used in Duan [31]. Garcia and Renault [48] showed that when the option price is homogeneous of degree one with respect to the asset price and the strike price, then the option delta hedge is consistent with Duan's formulation. Further detailed discussions on homogeneity of option prices and delta hedging for scale-invariant models are provided in Bates [11] and Alexander and Nogueira [1, 2].

The first approximation in (3.12) suggests that the minimum variance hedge is consistent with Duan's delta hedge when changes in today's asset values are positively correlated to changes in the next period's conditional variance. We further notice that the first specification for Δ_t depends on the probability measure under which the portfolio variance is minimized, while the second expression can be obtained under any probability space. A more detailed discussion on these two approximations and their relationship with the corresponding continuous time stochastic volatility limits for Gaussian driven GARCH models is provided in the Section 4.

The second part of Proposition 3.2 states that the local risk minimization hedge under a risk neutral measure Q converges to the total derivative expression from (3.12) when the time between two consecutive observations shrinks to zero. Moreover, the first approximation in **(i)** should also hold for the P - local risk minimization hedge, but only in the special case when the contingent claim price is computed based on the minimal martingale measure.

Huang and Guo [55] showed that the local risk minimization taken with respect to the risk neutral provided by the Extended Girsanov Principle converges to the Black Scholes delta hedge when the underlying is governed by a Geometric Brownian motion. This result can be viewed as a special case of **(ii)**.There are situations (see El Karoui *et al.* [39], Poulsen *et al.* [63]) in the continuous time setup and in the presence of a well defined minimal martingale measure, in which a volatility adjusted hedging ratio similar to (3.12) can be obtained out of local risk minimization, that is, without assumptions on the pricing kernel and by constructing a hedging portfolio that involves trading exclusively in the underlying asset and the riskless asset.

In the next two sections we shall investigate the empirical performance of the local risk minimization hedges and compare them with the instantaneous delta hedges that we described above for European call options.

4 Hedging Gaussian GARCH options

The goal of this section is to provide a theoretical and empirical analysis of the hedging schemes previously proposed for a general class of GARCH(1,1) models driven by Gaussian innovations. Consider then a GARCH(1,1) model as in (2.2) and (2.3) with a conditional mean prescription given by:

$$\mu_t := r + \lambda\sigma_t - \kappa^P_{\epsilon_t}(\sigma_t), \quad \lambda \in \mathbb{R}. \tag{4.1}$$

The main reason for choosing such a parametrization for the conditional mean return is that one can interpret λ as the market price of risk per unit volatility. Since, we also assume that the conditional distribution $\mathbf{D}(0,1)$ of the innovations is normal we have that $\kappa^P_{\epsilon_t}(\sigma_t) = \sigma_t^2/2$ and hence the return dynamics under the historical measure are identical to those in Duan [31], namely:

$$\begin{aligned} y_t &= r + \lambda\sigma_t - \frac{1}{2}\sigma_t^2 + \sigma_t\epsilon_t, \quad \epsilon_t \sim \mathbf{N}(0,1), & (4.2)\\ \sigma_t^2 &= \alpha_0 + \alpha_1\sigma_{t-1}^2\omega(\epsilon_{t-1}) + \beta_1\sigma_{t-1}^2. & (4.3)\end{aligned}$$

The Gaussian hypothesis makes identical the martingale measures Q_{egp} and Q_{ess} introduced in Section 2 so through this section we shall write just Q. Moreover, the specific conditional mean prescription chosen in (4.1) implies that the unit risk premium ϱ_t in (2.10) is constant and equal to λ, that is, $\varrho_t = \lambda$; this simplifies many of the computations that follow in this section. Then, under either of the martingale measures Q (that is, Q_{egp} or Q_{ess}) we can write:

$$\begin{aligned} y_t &= r - \frac{\sigma_t^2}{2} + \sigma_t\epsilon_t^*, \quad \epsilon_t^* \sim \mathbf{N}(0,1), & (4.4)\\ \sigma_t^2 &= \alpha_0 + \alpha_1\sigma_{t-1}^2\omega(\epsilon_{t-1}^* - \lambda) + \beta_1\sigma_{t-1}^2. & (4.5)\end{aligned}$$

This specification coincides to the one obtained by Duan [31] by applying the local risk neutral valuation principle. We now write down two expressions that we shall need in deriving the vega of a European call option that uses this GARCH model for the underlying asset. First, using (4.4)-(4.5), the time evolution of the prices under Q is given by

$$S_T = S_t e^{r(T-t)-\frac{1}{2}\sum_{l=t+1}^{T}\sigma_l^2+\sum_{l=t+1}^{T}\sigma_l\epsilon_l^*}. \tag{4.6}$$

Secondly, we let

$$P_i := \alpha_1 \omega(\epsilon_i^* - \lambda) + \beta_1, \quad i = 0, \ldots, T.$$

A straightforward inductive argument using (4.5) implies that

$$\sigma_s^2 = \alpha_0 A(s,l) + \sigma_l^2 B(s,l) \quad \text{with} \quad l = 0, \ldots T-1 \quad \text{and} \quad s > l,$$

and

$$A(s,l) := \prod_{i=l}^{s-1} P_i \left[\sum_{k=l}^{s-1} \frac{1}{\prod_{i=l}^{k} P_i} \right] \quad \text{and} \quad B(s,l) := \prod_{i=l}^{s-1} P_i. \tag{4.7}$$

The delta and vega of an European call option based on the Gaussian GARCH model are computed below.

Proposition 4.1 *Given a European call option with strike price K that has (4.2)-(4.3) as log-returns generating process, the corresponding delta and vega hedges are given by:*

$$\Delta_t^S = e^{-r(T-t)} E^Q \left[\frac{S_T}{S_t} \mathbf{1}_{\{S_T \geq K\}} \mid \mathcal{F}_t \right]. \tag{4.8}$$

$$\Delta_t^\sigma = \frac{e^{-r(T-t)}}{2} E^Q \left[S_T \left(\sum_{l=1}^{T-t} \left(\frac{\epsilon_{t+l}^*}{\sigma_{t+l}} - 1 \right) B(t+l, t+1) \right) \mathbf{1}_{\{S_T \geq K\}} \mid \mathcal{F}_t \right], \tag{4.9}$$

Here $B(\cdot,\cdot)$ is given as in (4.7) with the convention $B(t+1,t+1) := 1$.

Equation (4.8) resembles the formula derived in Duan [31] for Gaussian GARCH. We provide an alternative proof of this in the appendix and we argue that this relationship also holds for any type of asset return process with a risk-neutralization given as in (2.6)-(2.7). Formula (4.9) gives a representation for the option vega; a similar expression for the vega of a European call was given in Duan [35] for an asymmetric GARCH model, but without proof or numerical implementation. We provide a detailed proof for this expression in the appendix. Since there are no closed form results available for the conditional expectations in the thesis of Proposition 4.1, we use Monte Carlo simulations for evaluating them.

In order to compute the minimum variance hedge we only need to compute the correction terms in the two approximations in (3.12). We shall do so later by making specific assumptions about the form of the function $\omega(\cdot)$ in (4.3); at this stage, we only write down the expression for the derivative of the conditional variance with respect to the asset price since no requirements on the volatility specification are required. Indeed, since $S_t = S_{t-1} e^{r + \lambda\sigma_t - \frac{1}{2}\sigma_t^2 + \sigma_t \epsilon_t}$, we solve for ϵ_t and we obtain

$$\epsilon_t = \frac{1}{\sigma_t} \left[\log(S_t) - \log(S_{t-1}) - r - \lambda\sigma_t + \frac{1}{2}\sigma_t^2 \right].$$

If we substitute this expression into (4.3) we can easily compute the VM using the chain rule and obtain:

$$\frac{d\sigma_{t+1}^2}{dS_t} = \frac{\alpha_1 \sigma_t}{S_t} \frac{d\omega(\epsilon_t)}{d\epsilon_t}. \tag{4.10}$$

The minimum variance hedge is then obtained by substituting (4.8), (4.9) and (4.10) into (3.12). The empirical performance of this hedging ratio is tested in a numerical experiment at the end of this section. In the next subsection, we propose a different formula for the minimum variance hedge using an alternative approach based on a continuous time limit for the correction term used in the first approximation in (3.12).

4.1 Local risk minimization in stochastic volatility GARCH models

In this section we investigate the local risk minimization hedging strategy within the class of stochastic volatility models which are obtained as weak limits of Gaussian GARCH models and study its relationship to the minimum variance hedge that we derived in the previous section. The weak convergence of GARCH models to bivariate diffusions where the volatility process and the price process are uncorrelated was first studied by Nelson [61]. Following this approach Duan [33] studied the continuous time limit of a more general class of augmented GARCH models, which includes the most commonly used GARCH prescriptions in the literature. The convergence results are typically based on a set of non-unique assumptions on the model parameters. Non-uniqueness leads to the existence of other diffusion limits in the literature (see for example Corradi [29], Heston and Nandi [51] or Kluppelberg *et al.* [59] among others). Since the bivariate diffusion limits of GARCH models proposed by Nelson [61] and Duan [33] are of widespread use for pricing and hedging derivatives in the continuous time setting, we restrict our attention to those.

In order to emphasize the effect of the correlation between the the price process and the variance process on the minimum variance hedging, we investigate the continuous limit of a GARCH model that incorporates a leverage effect. More specifically, we consider the class of the asymmetric NGARCH models, which corresponds to choosing $\omega(\epsilon_t) = (\epsilon_t - \gamma)^2$ in (4.3), where γ is the leverage parameter. Following Duan [33], we first consider a finer discretization of the model from (4.2)-(4.3) by dividing the interval $[0, \ldots, T]$ into nT equally spaced points of length $h = 1/n$ and indexed by k, $k = 1, 2, \cdots, nT$. The IID sequence of standard Gaussian driving innovations is denoted by ϵ_k for any $k = 1, 2, \ldots, nT$. Let $\mathcal{F}_{n,k}$ be the information set generated by $\epsilon_1, \ldots, \epsilon_k$. Under these assumptions, the return and conditional variance have the following representation under the historical measure P:

$$\ln S_{k,h} - \ln S_{k-1,h} = \left(r + \lambda\sigma_{k,h} - \frac{1}{2}\sigma_{k,h}^2\right) h + \sigma_{k,h}\sqrt{h}\epsilon_k, \quad \epsilon_k \sim \mathbf{N}(0,1), \tag{4.11}$$

$$\sigma_{k+1,h}^2 - \sigma_{k,h}^2 = \left(\alpha_0 + \sigma_{k,h}^2(\beta + \alpha_1(1+\gamma^2) - 1)\right) h + \alpha_1\sigma_{k,h}^2\left((\epsilon_k - \gamma)^2 - (1+\gamma^2)\right)\sqrt{h}. \tag{4.12}$$

When $n = 1$, the above process resembles the standard NGARCH(1,1) specification of (4.2)-(4.3). This model can be further extend to a bivariate continuous time process as follows:

$$\left(S_{t,h}, \sigma_{t,h}^2\right) := \left(S_{k,h}, \sigma_{k+1,h}^2\right), \quad t \in [kh, (k+1)h), \;\; k = 0, 1, \ldots, nT - 1.$$

Duan (1997) [] showed that when h approaches zero, $\left(S_{t,h}, \sigma_{t,h}^2\right)$ converges weakly to a bivariate diffusion $\left(S_t, \sigma_t^2\right)$ which satisfies the following stochastic differential equations:

$$d\ln S_t = \left(r + \lambda\sigma_t - \frac{1}{2}\sigma_t^2\right) dt + \sigma_t dB_{1t}, \tag{4.13}$$

$$d\sigma_t^2 = \left(\alpha_0 + \sigma_t^2(\beta + \alpha_1(1+\gamma^2) - 1)\right) dt - 2\alpha_1\gamma\sigma_t^2 dB_{1t} + \sqrt{2}\alpha_1\sigma_t^2 dB_{2t}. \tag{4.14}$$

Here, B_{1t} and B_{2t} are two independent standard Brownian motions under P. Applying a discretized version of the extended Girsanov transform to the bivariate process given in (4.11)-(4.12), the risk neutral specifications become:

$$\ln S_{k,h} - \ln S_{k-1,h} = \left(r - \frac{1}{2}\sigma_{k,h}^2\right) h + \sigma_{k,h}\sqrt{h}\epsilon_k^*, \quad \epsilon_k^* \sim \mathbf{N}(0,1), \tag{4.15}$$

$$\sigma_{k+1,h}^2 - \sigma_{k,h}^2 = \left(\alpha_0 + \sigma_{k,h}^2(\beta + \alpha_1(1+\gamma^2) - 1)\right) h + \alpha_1\sigma_{k,h}^2\left((\epsilon_k^* - \gamma - \lambda\sqrt{h})^2 - (1+\gamma^2)\right)\sqrt{h}. \tag{4.16}$$

Taking the limit when $h \to 0$ of $\left(S_{t,h}, \sigma^2_{t,h}\right)$ under the martingale measure Q given by the Extended Girsanov Principle (or conditional Esscher transform), we get:

$$\begin{aligned} d\ln S_t &= \left(r - \frac{1}{2}\sigma_t^2\right) dt + \sigma_t dB^*_{1t}, && (4.17)\\ d\sigma_t^2 &= \left(\alpha_0 + \sigma_t^2(\beta + \alpha_1(1+\gamma^2) - 1 + 2\alpha_1\gamma\lambda)\right) dt - 2\alpha_1\gamma\sigma_t^2 dB^*_{1t} + \sqrt{2}\alpha_1\sigma_t^2 dB^*_{2t}. && (4.18)\end{aligned}$$

Here, B^*_{1t} and B^*_{2t} are two independent standard Brownian motions under Q, such that $dB^*_{1t} = dB_{1t} + \lambda dt$ and $dB^*_{2t} = dB_{2t}$. An interesting aspect is provided by the fact that equations (4.17)-(4.18) can be obtained directly by applying the continuous time version of the minimal martingale measure to the bivariate diffusion from (4.13)-(4.14). A more detailed discussion about this is provided in Duan [32]. In the next proposition we compute the local risk minimization hedging strategy for the continuous time limit of the NGARCH process.

Proposition 4.2 *Suppose that the asset returns are governed by the stochastic volatility process determined by (4.13)-(4.14). Then the price* Π^{sv}_t *and the local risk minimizing delta hedge* Δ^{sv}_t *of a European option with payoff* H *and maturity* T*, conditional on initial values* (x,y) *for the asset price and volatility process, respectively, are given by:*

$$\begin{aligned} \Pi^{sv}_t(x,y) &= e^{-r(T-t)} E^Q\left[H \mid \mathcal{F}^{x,y}_t\right], && (4.19)\\ \Delta^{sv}_t(x,y) &= \Delta^S_t(x,y) - 2\alpha_1\gamma\frac{y}{x}\Delta^\sigma_t(x,y). && (4.20)\end{aligned}$$

Here $\mathcal{F}^{x,y}_t = \mathcal{F}_t \cup \{(S_t,\sigma_t) = (x,y)\}$*, with* $\mathcal{F}_t$ *being the standard filtration in the stochastic volatility setup and* $\Delta^S_t(x,y)$ *and* $\Delta^\sigma_t(x,y)$ *are the standard option delta and vega conditional on the initial values* $S_t = x$ *and* $\sigma_t = y$ *for the asset price and the conditional volatility process. The option prices are computed based on the minimal martingale measure* Q *under which the asset returns process follow (4.17)-(4.18)*

The proof of Proposition 4.2 follows the same three steps procedure as in Poulsen *et al.* [63]; this algorithm follows the construction from El Karoui *et al.* [39] in which the market is first completed by introducing another tradable asset, then the delta hedge is computed in this setting and finally projected onto the original market. We notice two particular cases: first, $\alpha_1 = 0$ corresponds to a deterministic conditional variance and the local risk minimization hedging reduces to the standard Black Scholes delta hedge. Second, when $\gamma = 0$ the two Brownian motions are uncorrelated and in this situation the VM equals zero, so the option hedge corresponds to $\Delta^S_t(x,y)$.

Based on the discretization proposed above and its connection to the continuous time setting, we propose a different instantaneous delta hedge for the Gaussian GARCH setting. The next result serves as a justification for the use of that hedge instead of the one proposed by Duan [31] and shows that the VMs of the GARCH models converge to their continuous time counterparts.

Proposition 4.3 *Suppose that* $(S_{k,h}, \sigma^2_{k+1,h})$ *follows the dynamics of (4.11)-(4.12) under the historical measure* P *and (4.15)-(4.16) under the risk neutral measure* Q*. Then the following relation holds:*

$$\lim_{h\to 0} \frac{\mathrm{Cov}^P(S_{k+1,h} - S_{k,h}, \sigma^2_{k+2,h} - \sigma^2_{k+1,h} \mid \mathcal{F}^{x,y}_{k,h})}{\mathrm{Var}^P[S_{k+1,h} - S_{k,h} \mid \mathcal{F}^{x,y}_{k,h}]} = \lim_{h\to 0} \frac{\mathrm{Cov}^Q(S_{k+1,h} - S_{k,h}, \sigma^2_{k+2,h} - \sigma^2_{k+1,h} \mid \mathcal{F}^{x,y}_{k,h})}{\mathrm{Var}^Q[S_{k+1,h} - S_{k,h} \mid \mathcal{F}^{x,y}_{k,h}]} = -2\alpha_1\gamma\frac{y}{x} \quad (4.21)$$

Here $\mathcal{F}_{k,h}^{x,y} = \mathcal{F}_{k,h} \cup \{(S_{k,h}, \sigma_{k,h}) = (x,y)\}$, *with* $\mathcal{F}_{k,h}$ *being the standard filtration generated by the GARCH innovations.*

The proof is provided in the Appendix. Based on the continuous time analogy, we argue that the instantaneous local risk minimization delta hedge from (3.13) resembles Duan's delta hedge when either there are no ARCH effects or changes in asset prices are uncorrelated to changes in conditional variances and this translates to the absence of a leverage effect. When $\gamma \neq 0$ one should include a correction term into Duan's delta hedging formula. Proposition 4.3 indicates that, if the time between two consecutive trades shrinks to zero and the GARCH greeks converge to the corresponding stochastic volatility greeks, then the minimum variance hedge approaches the local risk minimization stochastic volatility hedge. Moreover, from (3.13), the Gaussian GARCH local risk minimization hedge under Q converges to its continuous time counterpart. In view of these arguments, we propose the following alternative delta hedge under the NGARCH model:

$$\Delta_t^{mvsv} = \Delta_t^S - 2\alpha_1\gamma\frac{\sigma_t}{S_t}\Delta_t^\sigma. \tag{4.22}$$

When fitted to financial return data, the asymmetric GARCH model typically yields a negative leverage effect, $\gamma > 0$, so a non-negative value of the option vega leads to a smaller value of the adjusted hedge ratio from (4.22) when compared to the one proposed by Duan [31]. In the numerical experiment from the next subsection, we implement this continuous time adjusted delta hedge, Δ_t^{mvsv}, as well as the second approximation from (3.12), which in the NGARCH case leads to:

$$\Delta_t^{mv} = \Delta_t^S + 2\alpha_1(\epsilon_t - \gamma)\frac{\sigma_t}{S_t}\Delta_t^\sigma \tag{4.23}$$

Although the results presented in this section are based on the NGARCH model, the same approach for the adjusted delta hedge can be implemented for other GARCH models by spelling out their continuous time limit.

4.2 Numerical comparisons of local risk minimization schemes under Gaussian GARCH models

In this section we provide numerical experiments to compare the hedging performance of our proposed schemes for the NGARCH(1,1) option pricing model from (4.2)-(4.3) with $\omega(\epsilon_t) = (\epsilon_t - \gamma)^2$. In order to work with realistic parameters, we estimate our model based on historical daily closing prices of the S&P 500 index from January 2, 1990 to August 30 2005, for a total of 3,951 observations. The NGARCH parameter values are consistent with what is usually found in the financial econometrics literature and are given below:

$$\alpha_0 = 1.1 \cdot 10^{-6}, \qquad \alpha_1 = -0.0556, \qquad \beta_1 = 0.8894, \qquad \lambda = 0.0302, \qquad \gamma = -0.9098.$$

We also assume the the annual interest rate is 0.7%. As in Ortega [62], for each hedging strategy considered, we compute the mean square hedging error ($MSHE$) under the physical probability as follows:

$$MSHE(\hat{\xi}) = E^P\left[\left(H - \hat{V}_0 - \sum_{k=1}^{T/j}\hat{\xi}_k^{(j)} \cdot (S_{kj} - S_{(k-1)j})\right)^2\right] \tag{4.24}$$

Since our study focuses on European Call options, the payoff at maturity of the option to hedge is $H = (S_T - K)_+$. $\hat{V}_0$ represents the option price and it is computed based on the NGARCH model with Gaussian innovations using a Monte Carlo simulation, $\hat{V}_0 = e^{-rT} E^Q \left[(S_T - K)_+ \right]$. The four competing hedging strategies $\hat{\xi}^{(j)}$ (the upper script (j) indicates that rebalancing takes place every j days) are summarized below:

- $\xi = (\xi_t)_{t \in \{j, 2j, \ldots, T\}}$, where ξ_t is the Q local risk minimization (LRM) hedge computed using equation (3.10).
- $\Delta^S = (\Delta_t^S)_{t \in \{0, j, \ldots, T-j\}}$, where Δ_t^S is the instantaneous delta hedge proposed by Duan [31] and computed using equation (4.8).
- $\Delta^{mv} = (\Delta_t^{mv})_{t \in \{0, j, \ldots, T-j\}}$, where Δ_t^{mv} is the minimum variance instantaneous delta hedge computed using equation (4.23).
- $\Delta^{mvsv} = (\Delta_t^{mvsv})_{t \in \{0, j, \ldots, T-j\}}$, where Δ_t^{mvsv} is the stochastic volatility adjusted minimum variance instantaneous delta hedge computed using equation (4.22).

Most empirical studies on GARCH option pricing and hedging include the Black-Scholes model into their comparison. There are two reasons we discard such a model from our simulation experiments: firstly, Ortega [62] found empirical evidence that both the local risk minimization and Duan's [31] outperform the Black-Scholes delta hedge, and secondly, since we prefer to investigate only the effect of the hedging scheme used in the behavior of the $MSHE$, it is better to put aside the Black-Scholes strategy because it uses an option price different to the GARCH one.

Since there are no closed form expressions available, all quantities necessary to compute the above hedging ratios require Monte Carlo simulations. As previously mentioned, this can be carried out in two ways: either by simulating the price paths directly under Q, or by simulating the paths under P and averaging the quantities of interest adjusted by the corresponding Radon-Nikodym derivatives. We adopt here the second approach since this leads to a variance reduction in the Monte Carlo estimators (see for example Badescu and Kulperger [3] for a discussion about computing option prices by simulating under the historical measure P).

We run three different exercises depending on the time to maturity T of our options and on the hedging frequency:

- **Exercise 1**: $T = 6$, 8, 10 and 12 days and rebalancing is done in a daily basis.
- **Exercise 2**: $T = 10$, 20, 30 and 40 days and rebalancing is done every 10 days.
- **Exercise 2**: $T = 20$, 40, 60 and 80 days and rebalancing is done every 20 days.

For all our simulations we consider the same starting value for the asset price, $S_0 = 100$. In order to investigate the sensitivity of the hedging method to the option moneyness, we consider three strike prices, $K = 90$, 100 and 110. We use the unconditional variance for the NGARCH model as the starting variance for our numerical simulations. The option prices, $\hat{V}_0$ are computed based on 10,000 random price paths and each expectation from the hedging

Table 4.1: Gaussian NGARCH average Mean Square Hedging Errors ($MSHE$) (computed across all maturities) for each moneyness and exercise. Average values for the corresponding standard errors are reported in the brackets.

	Moneyness	ξ	Δ^S	Δ^{mv}	Δ^{mvsv}
Exercise 1	1.11	0.4049	0.0136	0.0286	0.0209
		(0.0371)	(0.0029)	(0.0062)	(0.0045)
	1.00	0.2342	0.2159	0.2039	0.2074
		(0.0162)	(0.0159)	(0.0157)	(0.0144)
	0.9	0.0016	0.0022	0.0020	0.0014
		(0.0006)	(0.0009)	(0.0008)	(0.0007)
Exercise 2	1.11	0.5429	0.3853	0.4904	0.3645
		(0.0930)	(0.1049)	(0.1064)	(0.0818)
	1.00	1.4611	1.5502	1.4774	1.4026
		(0.1253)	(0.1685)	(0.1382)	(0.1211)
	0.9	0.1172	0.1602	0.1405	0.1139
		(0.0227)	(0.0202)	(0.0269)	(0.0239)
Exercise 3	1.11	1.8036	1.2805	1.9367	1.4446
		(0.3991)	(0.3574)	(0.4909)	(0.3770)
	1.00	3.1631	3.2265	3.5204	3.0039
		(0.3511)	(0.4067)	(0.4155)	(0.3444)
	0.9	0.9276	1.2082	1.2257	0.9147
		(0.2223)	(0.2042)	(0.2509)	(0.2184)

calculations are based on 5,000 Monte Carlo paths. The mean square hedging error associated with every hedging strategy will be computed by taking the mean of the terminal square hedging errors over 1000 price paths.

We start our analysis by examining the aggregate $MSHE$ for each exercise. The very short term options with a daily rebalancing indicate that all instantaneous delta hedges (Δ^S, Δ^{mv} and Δ^{mvsv}) produce almost the same $MSHE$ value of around 0.077, which is much lower than the corresponding $MSHE$ for the local risk minimization hedge of 0.215. For the second exercise, the overall performance of our methods is much closer, the stochastic volatility adjusted minimum variance hedge having a slightly lower error of around 0.63, all other methods being at 0.7. The longer term option combined with the lower frequency rebalancing suggest that Δ^{mvsv} ($MSHE = 1.78$) is the overall best strategy, while Δ^{mv} ($MSHE = 2.22$) being the worst. We can also conclude that the minimum variance approximation does not work very well in terms of this hedging criterion measure.

Additional interesting conclusions can be drawn by analyzing the hedging errors for each moneyness considered. The average values for the $MSHE$ (computed across maturities) and the corresponding standard error averages are presented in Table 4.1. All three exercises document that the behavior of our hedging scheme proposed depends crucially on the option's moneyness. For example, for the in-the-money (ITM) options in Exercises 1 and 3, we note that Duan's delta hedge provides the lowest $MSHE$, while its performance for Exercise 2 comes into a very close second place (Δ^{mvsv} is the best in this case). The out of-the-money (OTM) options consistently indicate that Δ^{mvsv} is the best method for all exercises, with the local risk minimizing strategy being also very close in value, ξ (the largest $MSHE$ difference between these two methods is 0.013. Both the other methods perform much worse in this case. For at-the-money (ATM) options there are no significant differences between the hedging strategies. However, we still note that Δ^{mvsv} represent the best scheme for Exercises 2 and 3, followed by ξ again. Moreover, the standard errors corresponding to these two methods are the smallest overall.

The smaller MSHE values for the deeper ITM and OTM European Call options confirm that these contracts are easier to be hedged compared to the ATM counterparts. Such findings have been also documented in the literature in various forms. For example, the recent study by Denkl *et al.* [30] which investigates the hedging performance of delta hedge ratios for exponential Levy processes, indicates a concave behaviour of the MSHE with respect to the spot price and for a fixed strike price. The MSHE increases as the option is near the money. The behaviour of the mean square is further illustrated in Figures 4.1. These graphs confirm our previous findings that Δ^{mvsv} and ξ outperform the other methods for longer term, ATM and OTM options, while the local risk minimization provide a poor alternative for very short maturity options hedged with daily frequency. The effect of the hedging horizon can be depicted by comparing the results for the same maturity and different exercise. For example, when $T = 40$, the hedging every 10 days instead of 20 days improves the mean square hedging error by 37% for Δ^S, 43% for ξ, 44% for Δ^{mv} and by 46% for Δ^{mvsv}. Thus, the proposed stochastic volatility adjusted delta hedge benefits the most when the hedging frequency is augmented.



(a) Exercise 1. The four maturities considered are 6, 8, 10, and 12 days and hedging is carried out daily.



(b) Exercise 2. The four maturities considered are 10, 20, 30, and 40 days and hedging is carried out every 10 days.



(c) Exercise 3. The four maturities considered are 20, 40, 60, 80 days and hedging is carried out every 20 days.

Figure 4.1: Mean square hedging error ($MSHE$) for the Gaussian NGARCH model and for the following hedging schemes, local risk

5 Numerical comparisons of Q-local risk minimization schemes under Normal Inverse Gaussian GARCH models

In this section we provide an empirical study of local risk minimization (LRM) hedging strategies constructed using a martingale measure for a special case of non-Gaussian GARCH models with Normal Inverse Gaussian (NIG) innovations. The NIG distribution has been studied in the context of the modeling of financial returns by Barndorff-Nielsen [8], and has been used in the ARCH setting by Jensen and Lunde [57]. More recently, the performance of GARCH option pricing models based on NIG innovations was investigated by Badescu *et al.* [6] for European style options and by Stentoft [68] for American options, among others. Empirical findings indicate that the NIG density is more appropriate for capturing the negative skewness and thicker tails generally displayed by asset returns than its Gaussian counterpart; moreover, option prices based on a NIG-GARCH model outperform the Gaussian benchmark. In the following paragraphs, we briefly review several notions related to the NIG distribution for the driving noise process and discuss its implication for option pricing based on the conditional Esscher transform and the Extended Girsanov Principle used in our numerical simulations.

We suppose that the innovations ϵ_t follow a conditional NIG distribution with parameters k, a, s and ℓ, that is, $\epsilon_t \mid \mathcal{F}_t \sim \mathbf{NIG}(k, a, s, \ell)$, and with probability density function given by:

$$f^P_{\epsilon_t}(x) = \frac{k}{\pi s} \exp\left[\sqrt{k^2 - a^2} + a\left(\frac{x-\ell}{s}\right)\right] \frac{K_1\left(k\sqrt{1 + \left(\frac{x-\ell}{s}\right)^2}\right)}{\sqrt{1 + \left(\frac{x-\ell}{s}\right)^2}}. \tag{5.1}$$

Here $K_1(\dot{})$ represents the modified Bessel function of third kind and index 1. The parameter k measures the kurtosis of the distribution, a the asymmetry, s is the scale parameter, and ℓ is the location of the distribution. Note that this NIG parametrization is particularly appropriate for practical implementations since it makes ϵ_t belong to a location-scale family of distributions; for this reason, k and a are called invariant parameters. The cumulant generating function is given by:

$$\kappa^P_{\epsilon_t}(z) = z\ell + \left(\sqrt{k^2 - a^2} - \sqrt{k^2 - (a + zs)^2}\right). \tag{5.2}$$

Thus, the first two conditional historical cumulants are:

$$\begin{aligned} E^P[\epsilon_t \mid \mathcal{F}_{t-1}] &= \frac{d\kappa^P_{\epsilon_t}(z)}{dz}\Big|_{z=0} = \ell + \frac{as}{\sqrt{k^2 - a^2}}, \\ \mathrm{Var}^P[\epsilon_t \mid \mathcal{F}_{t-1}] &= \frac{d^2\kappa^P_{\epsilon_t}(z)}{dz^2}\Big|_{z=0} = \frac{s^2 k^2}{(\sqrt{k^2 - a^2})^3}. \end{aligned}$$

Since the GARCH innovations have mean zero and unit variance this leads to a reduction in the NIG parameters which need to be estimated. The scale and the location can therefore be expressed in terms of the invariant parameters:

$$s = \frac{(\sqrt{k^2 - a^2})^{3/2}}{k}, \qquad \ell = -\frac{a}{k}(\sqrt{k^2 - a^2})^{1/2}\ .$$

From the scale-invariance property and (2.2) it follows that the asset returns are conditionally NIG distributed with the following parameters:

$$y_t \mid \mathcal{F}_t \sim \mathbf{NIG}(k, a, s\sigma_t, \mu_t + \ell\sigma_t). \tag{5.3}$$

We now have all the tools needed to derive risk-neutralized dynamics based on our NIG-GARCH model. Unlike the Gaussian case, the two measures proposed in Section 2 lead to different equations for the return process under the pricing measure. Although such dynamics are available and can be easily obtained by inserting the above quantities into Propositions 2.3 and 2.5, we undertake a similar approach as in the previous section by computing hedging errors based on the computation of cumulants under the historical measure P. This requires only the knowledge the P-returns dynamic and the two Radon-Nikodym derivatives corresponding to the pricing kernels. To fully specify the model, we assume the same structure for conditional mean return, μ_t as in (4.1) and the same asymmetric conditional variance specification. Thus, the historical dynamics for the returns is given by:

$$\begin{aligned} y_t &= r + \lambda\sigma_t - \kappa^P_{\epsilon_t}(\sigma_t) + \sigma_t\epsilon_t, \quad \epsilon_t \sim \mathbf{NIG}(k, a, s, \ell), && (5.4)\\ \sigma_t^2 &= \alpha_0 + \alpha_1\sigma_{t-1}^2(\epsilon_{t-1} - \gamma)^2 + \beta_1\sigma_{t-1}^2. && (5.5) \end{aligned}$$

Here the cumulant generating function is given in (5.2).

For the Extended Girsanov Principle, the market prices of risk is constant and the Radon-Nikodym derivative is given by:

$$\frac{dQ_{egp}}{dP} = \exp\left[\frac{a\lambda T}{s}\right]\prod_{t=1}^{T}\sqrt{\frac{s^2+(\epsilon_t-\ell)^2}{s^2+(\epsilon_t+\lambda-\ell)^2}}\,\frac{K_1\left(k\sqrt{1+\left(\frac{\epsilon_t+\lambda-\ell}{s}\right)^2}\right)}{K_1\left(k\sqrt{1+\left(\frac{\epsilon_t-\ell}{s}\right)^2}\right)}. \tag{5.6}$$

For the conditional Esscher transform, the Radon-Nikodym derivative is given by:

$$\frac{dQ_{ess}}{dP} = \exp\left[-T\sqrt{k^2-a^2}\right]\prod_{t=1}^{T}\exp\left[\hat{\theta}_t\sigma_t(\epsilon_t-\ell) + \sqrt{k^2-(a+\hat{\theta}_t\sigma_t s)^2}\right]. \tag{5.7}$$

For any $t \in [0, \cdots, T]$, the Esscher parameter $\hat{\theta}_t$ is the unique solution of the martingale equation (2.11). As previously mentioned, one of the advantages of using the NIG distribution in practice is that a closed form expression for $\hat{\theta}_t$ is available, namely:

$$\hat{\theta}_t = -\frac{1}{2} - \frac{a}{\sigma_t s} - \frac{1}{2}\sqrt{\frac{(\kappa^P_{\epsilon_t}(\sigma_t)-(\lambda+\ell)\sigma_t)^2}{\sigma_t^2 s^2}\left(\frac{4k^2}{\sigma_t^2 s^2 + (\kappa^P_{\epsilon_t}(\sigma_t)-(\lambda+\ell)\sigma_t)^2} - 1\right)}\,. \tag{5.8}$$

When an exact solution for $\hat{\theta}_t$ does not exist (for example the case of hyperbolic innovations) one can numerically solve (2.11) or use the second order Taylor approximation (2.12).

The numerical experiments that we present in this section are based on NIG-GARCH parameters obtained via maximum likelihood estimation (MLE) on the same data set of asset returns used in Section 4.2:

$$\alpha_0 = 8.8\cdot 10^{-7}, \qquad \alpha_1 = -0.0537, \qquad \beta_1 = 0.8946, \qquad \lambda = 0.0318, \qquad \gamma = -0.9157.$$

The estimated values for the NIG invariant parameters are, $k = 3.0897$ and $a = -0.3353$. As in the Gaussian case, we run the same three exercises that study different option maturities and rebalancing frequencies. Since one of the main objectives of this section is to examine the effect of the pricing measure in the hedging performance, we consider only local risk minimization (LRM), ξ, and Duan's (1995), Δ^S, as our hedging candidates. Both quantities are computed by simulating price paths under P and by making use of the Extended Girsanov Principle and the conditional Esscher transform via the corresponding Radon-Nikodym derivatives (5.6) and (5.7), respectively. The minimum variance delta hedges considered in the Gaussian NGARCH case are left out of this exercise because our continuous time limit diffusion approximation does not work in the case of the Esscher transform since the asset returns do not exhibit a standard NGARCH dynamic after the change of measure. The $MSHE$ is evaluated based on equation (4.24), the only difference being that the option price $\hat{V}_0$ is now computed separately for the each pricing measure. We use the same starting values for the asset price and the conditional starting variance, as well as the same number of Monte-Carlo simulations necessary for computing the prices and hedges as in the Gaussian case. The values for the $MSHE$ (computed across maturities) and the corresponding standard errors are presented in Table 5.2, while the dependencies of the $MSHEs$ with respect to each maturity are plotted in Figure 5.1 and Figure 5.2.

Table 5.2: NIG-NGARCH Mean Square Hedging Errors ($MSHE$) (computed across all maturities) for each moneyness and exercise. Values for the corresponding standard errors are reported between brackets.

	Moneyness	ξ_{egp}	Δ^S_{egp}	ξ_{ess}	Δ^S_{ess}
Exercise 1	1.11	0.4464	0.0196	0.4466	0.0197
		(0.0393)	(0.0055)	(0.0391)	(0.0055)
	1.00	0.3260	0.3088	0.3273	0.3102
		(0.0266)	(0.0267)	(0.0266)	(0.0267)
	0.9	0.0064	0.0088	0.0063	0.0088
		(0.0033)	(0.0047)	(0.0033)	(0.0047)
Exercise 2	1.11	0.6365	0.4031	0.6445	0.4023
		(0.0861)	(0.1095)	(0.0859)	(0.1089)
	1.00	1.4988	1.5911	1.5025	1.5949
		(0.1313)	(0.1645)	(0.1308)	(0.1639)
	0.9	0.1231	0.1994	0.1225	0.2017
		(0.0262)	(0.0421)	(0.0262)	(0.0424)
Exercise 3	1.11	1.8483	1.6573	1.8589	1.6538
		(0.2756)	(0.3821)	(0.2716)	(0.3796)
	1.00	3.5731	3.8464	3.5864	3.8684
		(0.3022)	(0.4289)	(0.3006)	(0.4298)
	0.9	0.9399	1.1849	0.9413	1.1983
		(0.1289)	(0.1164)	(0.1293)	(0.1171)

We first notice that there are no significant differences between the two pricing measures in terms of the hedging performance and their standard errors. Thus, we next look at the differences between the hedging prescriptions. Except for Exercise 1, for which LRM performs rather poorly, the overall average $MSHE$ of ξ_{egp}/ξ_{ess} and $\Delta^S_{egp}/\Delta^S_{ess}$ are almost equal. However, as in the Gaussian, case each method behaves differently depending on options' moneyness. For example, Duan's Δ outperforms the local risk minimization for all ITM options while the latter is the best hedging scheme for almost all other cases except ATM short term options (as in Exercise 1). A closer look at Table 4.1 and Table 5.2 for Exercises 2 and 3 reveals that the overall $MSHEs$ for the corresponding LRM strategies are not far from each other (with generally smaller standard errors for the NIG case). However, such a comparison may be misleading; indeed, the options prices that appear in (4.24) are different and moreover, they are computed together with the other necessary quantities via Monte Carlo simulations using different starting values of the conditional variance process. Therefore, in order to compare the hedging performance between GARCH models with different innovations, one has to take into account all the potential sources of errors which may end up playing an important role in the measure of the hedging performance.

6 Conclusions

In this paper we propose and analyze a variety of option hedging schemes for GARCH models driven by Gaussian and non-Gaussian innovations. Since the well-known minimal martingale measure is typically a signed measure for such models, the construction of local risk minimization (LRM) schemes with respect to the historical measure is not possible. Therefore, we first introduce the notion of LRM with respect to a martingale measure for general GARCH setups. The changes of measure used in our numerical experiments are the Extended Girsanov Principle and the conditional Esscher transform, for which we describe the risk neutral dynamics.

Next we analyze local risk minimization in the context of Gaussian GARCH models. We show that when time between two consecutive hedging trades is very small, the local risk minimization scheme can be approximated in two different ways. Both approximations can be viewed as generalizations of Duan's delta hedge formula [31] by taking into account the dependence of tomorrow's conditional variance on today's asset price. The first limit approximation is inspired in some of the existing results on the risk minimization for continuous time stochastic volatility models. More specifically, for a special asymmetric GARCH model, we propose a novel minimum variance delta hedge based on the continuous time bivariate diffusion limit of such a model. The second limit approximation is a minimum variance hedge that is obtained in a straightforward manner using chain rule differentiation. An extensive numerical experiment is constructed in order to compare these hedging schemes. Our simulation results suggest that the stochastic volatility inspired minimum variance hedge (Δ^{mvsv}) is the best overall hedging method in the sense that it provides the smallest mean square hedging error. The local risk minimization (ξ) comes into a close second place, although this method performs poorly for very short maturity options with high frequency rebalancing dates. After a more detailed data analysis, we found that both Δ^{mvsv} and ξ outperforms the other two methods for longer maturity options and lower hedging frequencies; interestingly, Duan's delta hedge is the best method for hedging out of-the-money options. Our results are consistent with some of the findings from the continuous time hedging literature according to which, local risk minimization provides the best hedge when the two driving Brownian motions of the corresponding stochastic volatility model are correlated.

Finally, we test the sensitivity of the local risk minimization and the Duan hedges with respect to the martingale measure used. Numerical experiments indicate there are no significant differences between the Extended Girsanov



(a) Exercise 1. The four maturities considered are 6, 8, 10, and 12 days and hedging is carried out daily.



(b) Exercise 2. The four maturities considered are 10, 20, 30, and 40 days and hedging is carried out every 10 days.



(c) Exercise 3. The four maturities considered are 20, 40, 60, 80 days and hedging is carried out every 20 days.

Figure 5.1: Mean square hedging error (MSHE) for the NIG-NGARCH model under the Extended Girsanov Principle for the schemes



(a) Exercise 1. The four maturities considered are 6, 8, 10, and 12 days and hedging is carried out daily.



(b) Exercise 2. The four maturities considered are 10, 20, 30, and 40 days and hedging is carried out every 10 days.



(c) Exercise 3. The four maturities considered are 20, 40, 60, 80 days and hedging is carried out every 20 days.

Figure 5.2: Mean square hedging error (MSHE) for the NIG-NGARCH model under the conditional Esscher transform for the schemes

Principle and the conditional Esscher transform. As in the Gaussian case, we found that the LRM outperforms Duan's delta hedge for long term options and lower hedging frequencies, while the latter is better for dealing with in-the-money options. It will be interesting to test and compare our hedging schemes for different types of Gaussian and non-Gaussian GARCH models on real option data sets, including very long term contracts. Since our stochastic volatility adjusted scheme outperformed all other candidates, other similar extensions can be proposed for a variety of non-Gaussian GARCH models.

References

[1] Alexander, C., and Nogueira, L. (2007). Model-free hedge ratios and scale-invariant models. *Journal of Banking and Finance*, 31(6), 1839-1861.

[2] Alexander, C., and Nogueira, L. (2007). Model-free price hedge ratios for homogeneous claims on tradable assets. *Quantitative Finance* 7(5), 473 479.

[3] Badescu, A.M., and Kulperger, R.J. (2008). GARCH option pricing: a semiparametric approach. *Insurance: Mathematics and Economics*, 43(1), 69-84.

[4] Badescu, A.M., Kulperger, R.J., and Lazar, E. (2008). Option valuation with Normal Mixture GARCH models, *Studies in Nonlinear Dynamics & Econometrics*, 12(2), Article 5.

[5] Badescu, A.M., Elliott, R.J., and Siu, T.K. (2009). Esscher transforms and consumption-based models, *Insurance: Mathematics and Economics*, 45(3), 337-347

[6] Badescu, A.M., Elliott, R.J., Kulperger, R.J., Miettinen, J. and Siu, T.K. (2011). A Comparison of Pricing Kernels for GARCH Option Pricing with Generalized Hyperbolic Distributions, *International Journal of Theoretical and Applied Finance*, 14(5), 669-708.

[7] Bakshi, G., Cao, C., and Chen, Z. (1997). Empirical performance of alternative option pricing models. *Journal of Finance*, 52, 2003-2049.

[8] Barndorff-Nielsen, O.E. (1997). Normal Inverse Gaussian Distributions and Stochastic Volatility Modelling. *Scandinavian Journal of Statistics*, 24(1), 1-13.

[9] Barone-Adesi, G., Engle, R., and Mancini, L. (2008). A GARCH Option Pricing Model in Incomplete Markets. *Review of Financial Studies*, 21, 1223-1258.

[10] Basak, S., and Chabakauri, G. (2012). Dynamic Hedging in Incomplete Markets: A Simple Solution. *Review of financial Studies*, 26(6), 1845-1896.

[11] Bates, D. (2005). Hedging the Smirk. *Finance Research Letters*, 2(4), 195-200.

[12] Black, F., and Scholes, M. (1973). The pricing of options and corporate liabilities. *Journal of Political Economy*, 81, 637-659.

[13] Bollerslev, T. (1986). Generalized Autoregressive Conditional Heteroskedasticity. *Journal of Econometrics*, 31, 307-327.

[14] Boyle, P., and Emanuel, D. (1980). Discretely adjusted option hedges. *Journal of Financial Economics*, 8, 259-282.

[15] Brennan, M.J. (1979). The pricing of contingent claims in discrete time models. *Journal of Finance*, 24(1), 53-68.

[16] Bühlmann, H., Delbaen, F., Embrechts, P., and Shiryaev, A.N. (1996). No-Arbitrage, change of measure and conditional Esscher transforms. *CWI Quarterly*, 9(4), 291-317.

[17] Černý, A. and Kallsen, J. (2007) On the structure of general mean-variance hedging strategies. *Annals of Probabability*, 35, 1479–1531.

[18] Černý, A. and Kallsen, J. (2008) A counterexample concerning the variance-optimal martingale measure. *Mathematical Finance*, 18, 305–316.

[19] Černý, A. and Kallsen, J. (2009) Hedging by sequential regressions revisited. *Mathematical Finance*, 19, 591–617.

[20] Chorro, C., Guegan, D. and Ielpo, F. (2011). Option Pricing for GARCH-type Models with Generalized Hyperbolic Innovations. *Quantitative Finance*, 12(7), 1079-1094.

[21] Christoffersen, P., and Jacobs, K. (2004). Which GARCH model for option valuation? *Management Science*, 50, 1204-1221.

[22] Christoffersen, P., Dorion, C., Jacobs, K., and Wang, Y. (2010). Volatility components: affine restrictions and Non-normal innovations. *Journal of Business and Economic Statistics.*, 28(4), 483-502.

[23] Christoffersen, P., Heston, S.L., and Jacobs, K. (2006). Option valuation with conditional skewness. *Journal of Econometrics*, 131, 253-284.

[24] Christoffersen, P., Heston, S.L., and Jacobs, K. (2012). Option Anomalies and the Stochastic Discount Factor. Working Paper, University of Maryland.

[25] Christoffersen, P., Redouane E., Feunou, B., and Jacobs, K. (2009) Option Valuation with Conditional Heteroskedasticity and Non-Normality. *Review of Financial Studies*, 23, 2139-2183.

[26] Christoffersen, P., Jacobs, K., and Ornthanalai, C. (2008). GARCH Option Valuation: Theory and Evidence. Working paper.

[27] Choi, Y. (2005). An analytical approximation to the option formula for the GARCH model, *International Review of Financial Analysis*, 14(2), 149-164.

[28] Cont, R., Tankov, P., and Voltchkova, E. (2005). Hedging with options in models with jumps. Abel Symposium 2005 on Stochastic analysis and applications in honor of Kiyosi Ito 90th birthday.

[29] Corradi, V. (2000). Reconsidering the Continuous Time Limit of the GARCH(1,1) Process. *Journal of Econometrics*, 96, 145-153.

[30] Denkl, S., Goy, M., Kallsen, J., Muhle-Karbe, J., and Pauwels, A. (2011). On the Performance of Delta Hedging Strategies in Exponential Levy Models, Working Paper.

[31] Duan, J.-C. (1995). The GARCH option pricing model. *Mathematical Finance*, 5, 13-32.

[32] Duan, J.-C., (1996). A unified theory of option pricing under stochastic volatility: from GARCH to diffusion. Working paper.

[33] Duan, J.-C. (1997) Augmented GARCH(p,q) processes and its diffusion limit. *Journal of Econometrics*, 79, 97-127.

[34] Duan, J.-C. (1999). Conditionally fat-tailed distributions and the volatility smile in options, unpublished manuscript, University of Toronto.

[35] Duan, J.-C. (2000). GARCH and Stochastic Volatility Option Pricing, unpublished manuscript.

[36] Duan, J.-C. and Simonato, J.-G. (1998) Empirical martingale simulation of asset prices. *Management Science*, 44(9), 1218-1233.

[37] Duan, J.-C., Gauthier, G., and Simonato, J.-G. (2001) Asymptotic distribution of the EMS option price estimator. *Management Science*, 47(8), 1122-1132.

[38] Duan, J.-C., Ritchken, P., and Sun, Z. (2006). Approximating GARCH-jump models, jump-diffusion processes, and option pricing. *Mathematical Finance*, 16(1), 21-52.

[39] El Karoui, N., Peng, S., and Quenez, M. C. (1997) Backward stochastic differential equations in finance. *Math. Finance*, 7, 1–71.

[40] Elliott, R.J., and Madan, D.G. (1998). A discrete time equivalent martingale measure. *Mathematical Finance*, 9(6), 825-841.

[41] Engle, R.F. (1982). Autoregressive conditional heteroskedasticity with estimates of the variance of U.K. inflation. *Econometrica*, 50, 987-1008.

[42] Engle, R.F., and Rosenberg, J. (1995). GARCH Gammas. *Journal of Derivatives*, 2, 47-59.

[43] Engle, R.F., and Rosenberg, J. (2000). Testing the Volatility Term Structure Using Option Hedging Criteria. *Journal of Derivatives*, 8(1), 10-28.

[44] Föllmer, H. and Schied, A. (2002) *Stochastic Finance*. Vol. 27 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co.

[45] Föllmer, H. and Sondermann, D. (1986) Hedging of nonredundant contingent claims. *Contributions to mathematical economics*, pp. 205–223, North-Holland.

[46] Föllmer, H. and Schweizer, M. (1989) Hedging by sequential regression: an introduction to the mathematics of option trading. *ASTIN Bull.*, 18, 147–160.

[47] Föllmer, H. and Schweizer, M. (1991) Hedging of contingent claims under incomplete information. *Applied stochastic analysis (London, 1989)*, vol. 5 of *Stochastics Monogr.*, pp. 389–414, Gordon and Breach.

[48] Garcia, R. and Renault, É. (1999) Note on Hedging in ARCH-type Option Pricing Models. *Mathematical Finance*, 8(2), 153-161.

[49] Gerber, H.U., and Shiu, E.S.W. (1994). Option pricing by Esscher transforms (with discussions). *Transactions of the Society of Actuaries* 46, 99-191.

[50] Gourieroux, C., and Monfort, A. (2007). Econometric specifications of stochastic discount factor models. *Journal of Econometrics*, 136, 509-530.

[51] Heston, S.L., and Nandi, S. (2000). A closed-form GARCH option pricing model. *Review of Financial Studies*, 13, 585-626.

[52] Hsieh, D.A. (1989). Modeling heteroskedasticity in daily foreign-exchange rates. *Journal of Business and Economic Statistics* 7, 307-317.

[53] Hsieh, K., and Ritchken, P. (2005). An empirical comparison of GARCH option pricing models. *Review of Derivative Research*, 8, 129-150.

[54] Huang, S.-F. and Guo, M. (2009) Financial derivative valuation—a dynamic semiparametric approach. *Statist. Sinica*, 19, 1037–1054.

[55] Huang, S.-F. and Guo, M. (2012) Dynamic programming and hedging strategies in discrete time. In Handbook of Computational Finance, (Edited by J. C. Duan, J. Gentle and W. Hardle), Springer, Berlin, 605-631.

[56] Hull, J., and White, A. (1987). The pricing of options on assets with stochastic volatility. *Journal of Finance*, 42, 281-300.

[57] Jensen, M. B., and Lunde, A. (2001). The NIG-S and ARCH model: a fat-tailed, stochastic, and autoregressive conditional heteroskedastic volatility model. *Econometrics Journal*, 4, 319-342.

[58] Kallsen, J. and Taqqu, M. S. (1998) Option pricing in ARCH-type models. *Mathematical Finance*, 8(1), 13-26.

[59] Kluppelberg, C., Lindner, A., and Maller, R. (2004). A continuous time GARCH process driven by Levy process: stationarity and second order behaviour. *Journal of Applied Probability*, 43 (3), 601- 622.

[60] Monfort, A. and Pegoraro, F. (2011). Asset pricing with second-order Esscher transforms. *The Journal of Banking and Finance*, 36, 1678-1687.

[61] Nelson, D. (1990). ARCH models as diffusion approximations. *Journal of Econometrics*, 45, 7-38.

[62] Ortega, J.-P. (2012). GARCH options via local risk minimization. *Quantitative Finance*, 12(7), 1095–1110.

[63] Poulsen, R., Schenk-Hoppé, K. R., and Ewald, C.-O. (2009). Risk minimization in stochastic volatility models: model risk and empirical performance. *Quant. Finance*, 9, 693–704.

[64] Prigent, J.L., and Scaillet, O. (2002). Weak Convergence of Hedging Strategies of Contingent Claims, Working paper.

[65] Rubinstein, M. (1976). The valuation of uncertain income streams and the pricing of options, *Bell Journal of Economics*, 7, 407-425.

[66] Schweizer, M. (2001) A guided tour through quadratic hedging approaches. *Option pricing, interest rates and risk management, Handbook of Mathematical Finance*, pp. 538–574, Cambridge University Press.

[67] Siu, T.K., Tong, H., and Yang, H. (2004). On pricing derivatives under GARCH models: a dynamic Gerber-Shiu approach. *North American Actuarial Journal*, 8(3), 17-32.

[68] Stentoft, L. (2008). American option pricing using GARCH models and the Normal Inverse Gaussian Distribution. *Journal of Financial Econometrics* 6, 540-582.

[69] Stentoft, L. and Rombouts, J. (2011). Option pricing with asymmetric heteroskedastic normal mixture models. Working paper.

[70] Wang, S.S. (2000). A class of distortion operators for pricing financial and insurance risks. *Journal of Risk and Insurance*, 67(1), 15-36.

7 Appendix

7.1 Proof of Proposition 2.3

(i) Using that ϱ_t is a $\mathcal{F}_t$-predictable process, we proceed by computing the one-step conditional mean of N_t:

$$E^P\left[N_t|\mathcal{F}_{t-1}\right] = \int \frac{f^P_{\epsilon_t}(x+\varrho_t)}{f^P_{\epsilon_t}(x)} f^P_{\epsilon_t}(x)dx = \int f^P_{\epsilon_t}(y)dy = 1.$$

Thus, the martingale property of Z_t follows imediatelly:

$$E^P\left[Z_t|\mathcal{F}_{t-1}\right] = Z_{t-1}E^P\left[N_t|\mathcal{F}_{t-1}\right] = Z_{t-1}.$$

Since $Z_T := dQ/dP > 0$ is by construction non-negative, the fact that Q is an equivalent probability measure with respect to P follows from:

$$E^Q[1] = E^P[Z_T] = E^P[Z_0] = 1.$$

(ii) Since $\epsilon^*_t := \epsilon_t + \varrho_t$, we notice that $\mathcal{F}_t = \sigma(\epsilon_1,\ldots,\epsilon_t) = \sigma(\epsilon^*_1,\ldots,\epsilon^*_t)$. We first show that ϵ_t under P and ϵ^*_t under Q are conditionally identically distributed by computing the corresponding cumulant generating functions assuming they exist. Let $u > 0$ and for any $z \in (-u,u)$ and using the Bayes rule, we have:

$$e^{\kappa^Q_{\epsilon^*_t}(z)} = E^Q\left[e^{z\epsilon^*_t}|\mathcal{F}_{t-1}\right] = E^P\left[e^{z(\epsilon_t+\varrho_t)}N_t|\mathcal{F}_{t-1}\right] = \int e^{z(x+\varrho_t)}\frac{f^P_{\epsilon_t}(x+\varrho_t)}{f^P_{\epsilon_t}(x)}f^P_{\epsilon_t}(x)dx = \int e^{zy}f^P_{\epsilon_t}(y)dy = e^{\kappa^P_{\epsilon_t}(z)}.$$

In particular, by taking the first and second order derivatives of the cumulant generating functions at zero, we notice that $E^Q\left[\epsilon^*_t|\mathcal{F}_{t-1}\right] = 0$ and $\mathrm{Var}^Q\left[\epsilon^*_t|\mathcal{F}_{t-1}\right] = 1$.

In order to show that ϵ^*_t are conditionally uncorrelated under Q we let $l > s > 1$. We have:

$$E^Q\left[\epsilon^*_t\epsilon^*_{t-s}|\mathcal{F}_{t-l}\right] = E^Q\left[E^Q\left[\epsilon^*_t\epsilon^*_{t-s}|\mathcal{F}_{t-1}\right]|\mathcal{F}_{t-l}\right] = E^Q\left[\epsilon^*_{t-s}E^Q\left[\epsilon^*_t|\mathcal{F}_{t-1}\right]|\mathcal{F}_{t-l}\right] = 0.$$

(iii) From equations (2.2) and (2.10) and using the previous result we have:

$$y_t = \mu_t + \sigma_t\epsilon_t = \mu_t + \sigma_t\left(\epsilon^*_t - \frac{\mu_t + \kappa^P_{\epsilon_t}(\sigma_t) - r}{\sigma_t}\right) = r - \kappa^Q_{\epsilon^*_t}(\sigma_t) + \sigma_t\epsilon^*_t.$$

The risk neutral property follows by verifying that the discounted asset prices satisfy the martingale property under Q. Indeed

$$E^Q\left[e^{y_t}|\mathcal{F}_{t-1}\right] = E^Q\left[e^{r-\kappa^Q_{\epsilon^*_t}(\sigma_t)+\sigma_t\epsilon^*_t}|\mathcal{F}_{t-1}\right] = e^{r-\kappa^Q_{\epsilon^*_t}(\sigma_t)}E^Q\left[e^{\sigma_t\epsilon^*_t}|\mathcal{F}_{t-1}\right] = e^r.$$

Replacing ϵ_t by $\epsilon^*_t - \varrho_t$ into the conditional variance equation (2.3) concludes the proof. ■

7.2 Proof of Proposition 2.5

(i) The proof follows the same steps as in the previous proposition after noticing that:

$$E^P\left[N_t|\mathcal{F}_{t-1}\right] = E^P\left[e^{\theta_t\sigma_t\epsilon_t-\kappa^P_{\epsilon_t}(\theta_t\sigma_t)}|\mathcal{F}_{t-1}\right] = 1.$$

(ii) Since $\epsilon_t^* := \frac{\epsilon_t - \kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t)}{\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}}$, the sigma algebras generated by the two innovation processes coincide. Next, we compute the conditional cumulant generating function of ϵ_t under Q. Let $u > 0$ and for any $z \in (-u, u)$ we have:

$$\begin{aligned}\kappa_{\epsilon_t}^Q(z) &= \ln E^Q\left[e^{z\epsilon_t}|\mathcal{F}_{t-1}\right] = \ln E^P\left[e^{z\epsilon_t}N_t|\mathcal{F}_{t-1}\right] = \ln E^P\left[e^{z\epsilon_t}e^{\theta_t\sigma_t\epsilon_t - \kappa_{\epsilon_t}^P(\theta_t\sigma_t)}|\mathcal{F}_{t-1}\right] \\ &= \kappa_{\epsilon_t}^P(z+\theta_t\sigma_t) - \kappa_{\epsilon_t}^P(\theta_t\sigma_t). \end{aligned} \tag{7.1}$$

Based on the above relation, we compute the first two conditional moments of ϵ_t by taking the first and second order derivatives of the cumulant generating function at the origin.

$$\begin{aligned} E^Q\left[\epsilon_t|\mathcal{F}_{t-1}\right] &= \left.\frac{d\kappa_{\epsilon_t}^Q(z)}{dz}\right|_{z=0} = \kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t) \\ \mathrm{Var}^Q\left[\epsilon_t|\mathcal{F}_{t-1}\right] &= \left.\frac{d^2\kappa_{\epsilon_t}^Q(z)}{dz^2}\right|_{z=0} = \kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t). \end{aligned}$$

Thus, using the definition of ϵ_t^*, we obtain immediately that $E^Q\left[\epsilon_t^*|\mathcal{F}_{t-1}\right] = 0$ and $\mathrm{Var}^Q\left[\epsilon_t^*|\mathcal{F}_{t-1}\right] = 1$. The fact that ϵ_t^* are conditionally uncorrelated follows in a similar way as in the previous proof.

(iii) From equation (2.2) it follows that:

$$y_t = \mu_t + \sigma_t\epsilon_t = \mu_t + \sigma_t\kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t) + \sigma_t\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}\epsilon_t^*. \tag{7.2}$$

If we denote $\sigma_t^* = \sigma_t\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}$, we only need to show that:

$$\mu_t + \sigma_t\kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t) = r - \kappa_{\epsilon_t^*}^Q(\sigma_t^*), \tag{7.3}$$

in order to obtained the required risk neutralized return dynamic under the conditional Esscher transform. We start by evaluation the cumulant generating function of ϵ_t^* under Q.

$$\kappa_{\epsilon_t^*}^Q(z) = \ln E^Q\left[e^{z\left(\frac{\epsilon_t - \kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t)}{\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}}\right)}\Big|\mathcal{F}_{t-1}\right] = -z\frac{\kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t)}{\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}} + \kappa_{\epsilon_t}^Q\left(\frac{z}{\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}}\right)$$

Using the above relation, (2.11), and (7.1) we have:

$$\begin{aligned} r - \kappa_{\epsilon_t^*}^Q(\sigma_t^*) &= r - \kappa_{\epsilon_t}^Q(\sigma_t) + \sigma_t\kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t) = r - \left(\kappa_{\epsilon_t}^P(\sigma_t+\theta_t\sigma_t) - \kappa_{\epsilon_t}^P(\theta_t\sigma_t)\right) + \sigma_t\kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t) \\ &= \mu_t + \sigma_t\kappa_{\epsilon_t}^{P\prime}(\theta_t\sigma_t), \end{aligned}$$

which proves (7.3). The risk neutral variance equation under the Esscher transform is obtained by multiplying the physical conditional variance equation (2.3) by $\sqrt{\kappa_{\epsilon_t}^{P\prime\prime}(\theta_t\sigma_t)}$. ■

7.3 Proof of Proposition 3.2

(i) The value X_t at a given time t of the hedge portfolio constructed in the statement is given by:

$$X = \Pi_t - \Delta_t^{mv}S_t.$$

If the hedge is kept constant over the next period, the change in value of the portfolio between two consecutive time steps is:

$$X_{t+1} - X_t = \Pi_{t+1} - \Pi_t - \Delta_t^{mv}\left(S_{t+1} - S_t\right). \tag{7.4}$$

When the time between the observations is small, we can use the following approximation for the change in the option price:

$$\Pi_{t+1} - \Pi_t \approx \frac{\partial \Pi_t}{\partial S_t} \left(S_{t+1} - S_t\right) + \frac{\partial \Pi_t}{\partial \sigma_{t+1}^2} \left(\sigma_{t+2}^2 - \sigma_{t+1}^2\right). \tag{7.5}$$

It follows that the one-period variance to be minimized is approximately given by:

$$\begin{aligned}
\operatorname{Var}^P\left(X_{t+1} - X_t \mid \mathcal{F}_t\right) &\approx \left(\frac{\partial \Pi_t}{\partial S_t} - \Delta_t^{mv}\right)^2 \operatorname{Var}^P\left[S_{t+1} - S_t \mid \mathcal{F}_t\right] \\
&+ 2\left(\frac{\partial \Pi_t}{\partial S_t} - \Delta_t^{mv}\right) \frac{\partial \Pi_t}{\partial \sigma_{t+1}^2} \operatorname{Cov}^P(S_{t+1} - S_t, \sigma_{t+2}^2 - \sigma_{t+1}^2 \mid \mathcal{F}_t) \\
&+ \left(\frac{\partial \Pi_t}{\partial \sigma_{t+1}^2}\right)^2 \operatorname{Var}^P\left[\sigma_{t+2}^2 - \sigma_{t+1}^2 \mid \mathcal{F}_t\right].
\end{aligned}$$

Taking the derivative of the above expression with respect to Δ_t^{mv} and setting it equal to zero, we have:

$$\Delta_t^{mv} \approx \Delta_t^S + \Delta_t^\sigma \frac{\operatorname{Cov}^P(S_{t+1} - S_t, \sigma_{t+2}^2 - \sigma_{t+1}^2 \mid \mathcal{F}_t)}{\operatorname{Var}^P[S_{t+1} - S_t \mid \mathcal{F}_t]}. \tag{7.6}$$

Moreover, if we further approximate the next period change in the conditional variance as

$$\sigma_{t+2}^2 - \sigma_{t+1}^2 \approx \frac{d\sigma_{t+1}^2}{dS_t} \left(S_{t+1} - S_t\right), \tag{7.7}$$

and plug it into (7.6) we obtain the second approximation for the minimum variance delta hedge.

(ii) When the local risk minimization is done with respect the risk neutral measure Q, we have $V_t = \Pi_t$. Thus, plugging in the approximations from (7.5) and (7.7) into the formula for the local risk minimization hedge under Q, we have:

$$\begin{aligned}
\xi_{t+1} &= \frac{\operatorname{Cov}^Q\left(\widetilde{V}_{t+1}, \widetilde{S}_{t+1} - \widetilde{S}_t \mid \mathcal{F}_t\right)}{\operatorname{Var}^Q\left[\widetilde{S}_{t+1} - \widetilde{S}_t \mid \mathcal{F}_t\right]} = \frac{\operatorname{Cov}^Q\left(V_{t+1}, S_{t+1} \mid \mathcal{F}_t\right)}{\operatorname{Var}^Q\left[S_{t+1} \mid \mathcal{F}_t\right]} \\
&\approx \frac{\operatorname{Cov}^Q\left(V_t + \frac{\partial \Pi_t}{\partial S_t}\left(S_{t+1} - S_t\right) + \frac{\partial \Pi_t}{\partial \sigma_{t+1}^2}\left(\sigma_{t+2}^2 - \sigma_{t+1}^2\right), S_{t+1} \mid \mathcal{F}_t\right)}{\operatorname{Var}^Q\left[S_{t+1} \mid \mathcal{F}_t\right]} \\
&= \Delta_t^S + \Delta_t^\sigma \frac{\operatorname{Cov}^Q(S_{t+1} - S_t, \sigma_{t+2}^2 - \sigma_{t+1}^2 \mid \mathcal{F}_t)}{\operatorname{Var}^Q[S_{t+1} - S_t \mid \mathcal{F}_t]} \\
&\approx \frac{\operatorname{Cov}^Q\left(V_t + \left(\frac{\partial \Pi_t}{\partial S_t} + \frac{\partial \Pi_t}{\partial \sigma_{t+1}^2} \frac{d\sigma_{t+1}^2}{dS_t}\right)\left(S_{t+1} - S_t\right), S_{t+1} \mid \mathcal{F}_t\right)}{\operatorname{Var}^Q\left[S_{t+1} \mid \mathcal{F}_t\right]} \\
&= \Delta_t^S + \Delta_t^\sigma \frac{d\sigma_{t+1}^2}{dS_t}. \quad \blacksquare
\end{aligned}$$

7.4 Proof of Proposition 4.1

We compute separately each term in (3.12) in our setup. Firstly, for any fixed t, we assume that certain regularity conditions which allow the interchange between the derivative and the expectation are satisfied. Using (4.6), we can write

$$\begin{aligned}\Delta_t^S &= \frac{\partial C}{\partial S_t}\left(S_t,\sigma_{t+1}^2(S_t)\right)=e^{-r(T-t)}\frac{\partial}{\partial S_t}E_t^Q\left[\max\left\{(S_T-K),0\right\}\right]\\ &= e^{-r(T-t)}\frac{\partial}{\partial S_t}E_t^Q\left[(S_T-K)\mathbf{1}_{\{S_T\geq K\}}\right]=e^{-r(T-t)}E_t^Q\left[\frac{\partial S_T}{\partial S_t}\mathbf{1}_{\{S_T\geq K\}}\right]\\ &= e^{-r(T-t)}E_t^Q\left[e^{r(T-t)-\frac{1}{2}\sum_{l=t+1}^{T}\sigma_l^2+\sum_{l=t+1}^{T}\sigma_l\epsilon_l^*}\mathbf{1}_{\{S_T\geq K\}}\right]=e^{-r(T-t)}E_t^Q\left[\frac{S_T}{S_t}\mathbf{1}_{\{S_T\geq K\}}\right]. \qquad (7.8)\end{aligned}$$

Second, we compute the option vega $\Delta_t^\sigma=\frac{\partial C}{\partial\sigma_{t+1}^2}\left(S_t,\sigma_{t+1}^2(S_t)\right)$. We start by using (4.7) in order to rewrite (4.6) as

$$S_T=S_t\left[e^{r(T-t)-\frac{1}{2}\sigma_{t+1}^2+\sigma_{t+1}\epsilon_{t+1}^*-\frac{1}{2}\left[\sum_{l=2}^{T-t}\alpha_0A(t+l,t+1)+\sigma_{t+1}^2B(t+l,t+1)\right]+\sum_{l=2}^{T-t}\left[\alpha_0A(t+l,t+1)+\sigma_{t+1}^2B(t+l,t+1)\right]^{\frac{1}{2}}\epsilon_{t+l}^*}\right].$$

Using this expression it is very easy to compute the derivative:

$$\frac{\partial S_T}{\partial\sigma_{t+1}^2}=\frac{S_T}{2}\left[\left(\frac{\epsilon_1^*}{\sigma_{t+1}}-1\right)+\sum_{l=2}^{T-t}\left(\frac{\epsilon_{t+l}^*}{\sigma_{t+l}}-1\right)B(t+l,t+1)\right]=\frac{S_T}{2}\left[\sum_{l=1}^{T-t}\left(\frac{\epsilon_{t+l}^*}{\sigma_{t+l}}-1\right)B(t+l,t+1)\right], \qquad (7.9)$$

where in the last equality we use the convention $B(t+1,t+1):=1$. We now use (7.9) to compute

$$\begin{aligned}\Delta_t^\sigma &= \frac{\partial C}{\partial\sigma_{t+1}^2}\left(S_t,\sigma_{t+1}^2(S_t)\right)=e^{-r(T-t)}\frac{\partial}{\partial\sigma_{t+1}^2}E_t^Q\left[\max\left\{(S_T-K),0\right\}\right]=e^{-r(T-t)}E_t^Q\left[\frac{\partial S_T}{\partial\sigma_{t+1}^2}\mathbf{1}_{\{S_T\geq K\}}\right]\\ &= \frac{e^{-r(T-t)}}{2}E_t^Q\left[S_T\left(\sum_{l=1}^{T-t}\left(\frac{\epsilon_{t+l}^*}{\sigma_{t+l}}-1\right)B(t+l,t+1)\right)\mathbf{1}_{\{S_T\geq K\}}\right]. \quad\blacksquare \qquad (7.10)\end{aligned}$$

7.5 Proof of Proposition 4.3

We only show the first equality for the second one follows in a similar fashion. Let $\sigma_{k+1,h}^2(x,y)$ and $\epsilon_k(x,y)$ be the restrictions of $\sigma_{k+1,h}^2$ and ϵ_k to the filtration $\mathcal{F}_{k,h}^{x,y}$. From (4.12) it follows that

$$\sigma_{k+1,h}^2(x,y)=y^2+a_1(y)h+a_2\left(y,\epsilon_k(x,y)\right)\sqrt{h}.$$

Here the functions a_1 and a_2 are given by:

$$\begin{aligned}a_1(y) &= \alpha_0+y^2(\beta+\alpha_1(1+\gamma^2)-1),\\ a_2(y,\epsilon_k(x,y)) &= \alpha_1y^2\left((\epsilon_k(x,y)+\gamma)^2-(1+\gamma^2)\right).\end{aligned}$$

It follows immediately that $\lim_{h\to0}\sigma_{k+1,h}(x,y)=y$. First, we compute the denominator of the left hand side of (4.21) for the Gaussian NGARCH model using 2.5. We have:

$$\begin{aligned}\mathrm{Var}^P\left[S_{k+1,h}-S_{k,h}\mid\mathcal{F}_{k,h}^{x,y}\right] &= x^2e^{2\left(r+\lambda\sigma_{k+1,h}(x,y)-\frac{1}{2}\sigma_{k+1,h}^2(x,y)\right)h}\mathrm{Var}^P\left[e^{\sigma_{k+1,h}\sqrt{h}\epsilon_{k+1}}\mid\mathcal{F}_{k,h}^{x,y}\right]\\ &= x^2e^{2\left(r+\lambda\sigma_{k+1,h}(x,y)-\frac{1}{2}\sigma_{k+1,h}^2(x,y)\right)h}\left(e^{\kappa_k(2\sigma_{k+1,h}(x,y)\sqrt{h})}-e^{2\kappa_k(\sigma_{k+1,h}(x,y)\sqrt{h})}\right)\\ &= x^2e^{2(r+\lambda\sigma_{k+1,h}(x,y))h}\left(e^{\sigma_{k+1,h}^2(x,y)h}-1\right).\end{aligned}$$

The conditional covariance between changes in asset price and conditional variances is given by:

$$\begin{aligned}\mathrm{Cov}^P\left(S_{k+1,h}-S_{k,h},\sigma_{k+2,h}^2-\sigma_{k+1,h}^2\mid\mathcal{F}_{k,h}^{x,y}\right) &= \mathrm{Cov}^P\left(S_{k+1,h},\sigma_{k+2,h}^2\mid\mathcal{F}_{k,h}^{x,y}\right)\\ &= x\alpha_1h\sqrt{h}\sigma_{k+1,h}^2(x,y)e^{\left(r+\lambda\sigma_{k+1,h}(x,y)-\frac{1}{2}\sigma_{k+1,h}^2(x,y)\right)h}\\ &\times\ \mathrm{Cov}^P\left(e^{\sqrt{h}\sigma_{k+1,h}\epsilon_{k+1}},(\epsilon_{k+1}+\gamma)^2\mid\mathcal{F}_{k,h}^{x,y}\right).\end{aligned}$$

Using the fact that $(\epsilon_{k+1}+\gamma)^2$ has a noncentral chi-squared distribution with one degree of freedom and noncentrality parameter γ^2, it follows that:

$$\mathrm{Cov}^P\left(e^{\sqrt{h}\sigma_{k+1,h}\epsilon_{k+1}}, (\epsilon_{k+1}+\gamma)^2 \mid \mathcal{F}_{k,h}^{x,y}\right) = e^{\frac{1}{2}\sigma_{k+1}^2(x,y)h}\sigma_{k+1,h}(x,y)\sqrt{h}\left(\sigma_{k+1,h}(x,y)\sqrt{h}+2\gamma\right),$$

and therefore, the numerator becomes:

$$\mathrm{Cov}^P\left(S_{k+1,h}-S_{k,h}, \sigma_{k+2,h}^2-\sigma_{k+1,h}^2 \mid \mathcal{F}_{k,h}^{x,y}\right) = x\alpha_1 h e^{(r+\lambda\sigma_{k+1,h}(x,y))h}\sigma_{k+1,h}^3(x,y)\left(\sigma_{k+1,h}(x,y)\sqrt{h}+2\gamma\right).$$

Taking the limit as h approaches zero we obtain:

$$\lim_{h\to 0}\frac{\mathrm{Cov}^P(S_{k+1,h}-S_{k,h}, \sigma_{k+2,h}^2-\sigma_{k+1,h}^2 \mid \mathcal{F}_{k,h}^{x,y})}{\mathrm{Var}^P[S_{k+1,h}-S_{k,h} \mid \mathcal{F}_{k,h}^{x,y}]} = \lim_{h\to 0}\frac{\alpha_1 h\sigma_{k+1,h}^3(x,y)\left(\sigma_{k+1,h}(x,y)\sqrt{h}+2\gamma\right)}{x\left(e^{\sigma_{k+1,h}^2(x,y)h}-1\right)} = 2\alpha_1\gamma\frac{y}{x}. \quad \blacksquare$$